

A case of melanism in a population of the insular lizard *Podarcis hispanica atrata*

Aurora M. CASTILLA

SHNB



SOCIETAT D'HISTÒRIA
NATURAL DE LES BALEARS

Castilla, A.M., 1994. A case of melanism in a population of the insular lizard *Podarcis hispanica atrata*. *Boll. Soc. Hist. Nat. Balears*, 37: 175-180 ISSN, 0212-260-X. Palma de Mallorca.

In the essay the finding and some morphological characteristics of a melanistic example of *Podarcis hispanica atrata* found on the Columbretes archipelago (Castellón, Spain) is described.

Key words: Mediterranean islands, lizard, melanism, endangered, endemism, age.

UN CASO DE MELANISMO EN UNA POBLACIÓ INSULAR DE LA LAGARTIJA. En este estudio se describe el hallazgo y algunas características morfológicas de un ejemplar melánico de *Podarcis hispanica atrata* encontrado en el archipiélago de las islas Columbretes (Castellón, España).

Palabras clave: Islas mediterráneas, lagartija, melanismo, en peligro, endemismo, edad.

Aurora M. CASTILLA. Instituto de Estudios Avanzados de las Islas Baleares (CSIC), Ctra. Valldemossa, Km. 7,500, E-07071 Palma de Mallorca, Spain, and Department of Biology, University of Antwerp (UIA), B-2610 Wilrijk, Belgium.

Recepció del manuscrit, 30-juny-94. Revisió acceptada, 11-oct-94

In this paper I describe the finding and some morphological and behavioural characteristics of a melanistic individual of the lizard *Podarcis hispanica atrata*. This is an endangered lizard whose world range is restricted to the Columbretes islands (39° 54' N, 0° 41' E) (see details in Castilla & Bauwens, 1991b). These constitute an archipelago of small uninhabited islets of volcanic origin situated in the Mediterra-

nean ca. 57 km off the coast of Castellón (province of Castellón, Spain). Both males and females show a great individual variability in the ventral colour and dorsal colour pattern. The latter ranges from pale uniform green to brownish-green with black reticulate spots (Castilla, L. Bauwens, in press).

This subspecies of lizard was described by Boscá (1916) on the base of a single individual that was brought

to him by a fisherman as a curiosity. The subspecific name *atrata* (Latin = dressed in black) refers to the black colour of that individual (Boscá, 1916). Since that date, however, no other melanistic individual has been seen or collected on the archipelago by herpetologists who made short-term visits to the islands in different periods (Eisentraut, 1930; Klemmer, 1961).

During the years 1984-1994 I have performed detailed studies of the lizards on Columbretes (see Castilla *et al.*, 1987; Castilla & Bauwens, 1991a, 1991b; Castilla & Swallow, in press), during which I handled several hundreds of individuals, and closely observed many others. Despite my intensive sampling efforts, only a single melanistic individual was caught in May 1991 on the northern and highest (ca. 70 m) part of the island Columbrete Grande (Figure 1). Thus, the occurrence of melanism seems to be an extremely rare event in *P. hispanica atrata*.

The melanistic individual was an adult male that had a coal bright uniform black colour, only the belly presented a black reddish coloration. This black coloration was darker than the black-greyish colour described for a melanistic individual of *Podarcis h. hispanica* found in the Iberian Peninsula (Pérez-Mellado, 1984). The single melanistic individual was observed throughout 1991-1994 and did not change his uniform bright black coloration. Hence, this must not be considered as a case of temporary melanism as described for turtles (Lovich *et al.*, 1990) and other vertebrates (see refs. in Sage, 1962), but instead, as a type of permanent melanism as defined in Lovich *et al.* (1990).

The body size (snout-vent length = 68.9 mm) and weight (8.86 g) of the melanistic individual was above the average recorded for adult male lizards in that population and year (Castilla & Bauwens, in press). In other species of reptiles, melanistic males attain a larger body size than normally coloured individuals (Lovich *et al.*, 1990). Upon its first capture, I estimated that the melanistic individual was at least two years old. The lizard was maintained temporarily in captivity during 1991. Afterwards, it was observed in the study area until October 1994, thus attaining an age of at least five years. No precise information is available on the age of other individuals from Columbretes. Nevertheless, data from a Portuguese population of *P. h. hispanica* indicate that the maximal age did not exceed three years (Caetano *et al.*, 1986). I suggest that the large body size of the melanistic lizard, and the reduced predation pressure on Columbretes (see in Castilla and Bauwens, 1991a,b) may have contributed to the advanced age (at least 5 years) attained by this individual.

The rare occurrence of melanistic individuals has been observed in continental populations of other lacertid lizards. Some examples are *P. hispanica* (Pérez-Mellado, 1984) and *P. muralis* (Zuffi, 1986; Barbadillo and Sánchez-Herráinz, 1992). It has been suggested that melanism is relatively common at northern latitudes (Kuranova, 1989), as for example in the lacertids *P. muralis* (Bruno and Maugeri, 1971), *Lacerta vivipara* (Malkmus, 1976), and *L. agilis*. On the other hand, some insular populations of mediterranean lizards (e.g. *P. dugesii*, *P. lilfordi*, *P. pityusen-*



Fig. 1. Melanistic male of *Podarcis hispanica atrata* in the Mediterranean island Columbrete Grande (a) (A. M. Castilla) and detail of the head (b) (L.J. Barbadillo).

Fig. 1. Macho melánico de Podarcis hispanica atrata en la isla Mediterránea de Columbrete Grande (a) (A. M. Castilla) y detalle de la cabeza (b) (L. J. Barbadillo).

sis) consist entirely of melanistic individuals (Bischoff *et al.*, 1989; Pérez-Mellado, 1989; Cirer and Martínez Rica, 1990). In general, it has been suggested that conditions associated with insularity, either on oceanic islands or on restricted alpine habitats, could favour the apparition of melanistic forms.

Acknowledgements

Permission to work in the island and transport to it was made possible by the Conselleria D'Medi Ambient (Generalitat Valenciana), the Dirección General de Puertos of Castellón, and the Jefatura Provincial de Aduanas of Castellón. Valentín Tena, Javier Martínez, Antonio Sanchez and Santiago Sales provided help during observations. My studies on Columbretes were funded by a grant of the Spanish M.E.C., the Project DGICYT (PB91-0055), and the Exmo. Ayuntamiento de Castellón.

References

- Barbadillo, L. J. & Sánchez-Herráiz, M. J. 1992. Melanismo en una población de *Podarcis muralis* (Reptilia, lacertidae) de Cantabria (N.de España). *Bol. Asoc. Herp. Esp.* 3: 15-17.
- Bischoff, W., Osenegg, K. & Mayer, W. 1989. Untersuchungen zur subspezifischen Gliederung der Madeira-Mauereidechse, *Podarcis dugesii* (Milne-Edwards 1829). *Salamandra*. 25: 237-259.
- Boscá, E. 1916. Dos observaciones a propósito de la *Lacerta muralis* en España. *Bol. R. Soc. Española Hist. Nat.* 16: 327- 330.
- Bruno, S. & Maugeri, S. 1977. *Rettili d'Italia*. Tartarughe- Sauri-Serpenti. Giunti Martello, Firenze.
- Caetano, M. H., Castanet, J. & Crespo, E. G. 1986. Estimation a l'aide de la squelettochronologie de l'âge de *Podarcis hispanica* (Steindachner, 1870), Sauria, Lacertidae, provenant d'une population portugaise. *Revue suisse Zool.* 93:117-127.
- Castilla, A. M. & Bauwens, D. 1991a. Thermal biology, microhabitat selection, and conservation of the insular lizard *Podarcis hispanica atrata*. *Oecologia*. 85: 366-374.
- Castilla, A. M. & Bauwens, D. 1991b. Observations on the natural history, present status, and conservation of the insular lizard *Podarcis hispanica atrata* on the Columbretes archipelago, Spain. *Biologia, Conservación de la lagartija Podarcis hispanica atrata en las islas Columbretes*. Ed. Generalitat Valenciana Conselleria de Medio Ambiente, Valencia 58: 69-84.
- Castilla, A. M. & J. Swallow. in press. Artificial egg-laying sites for lizards: A conservation strategy. *Biol. Conserv.*
- Cirer, A. M. & Martínez-Rica, J. P. 1990. The polymorphism of *Podarcis pityusensis* and its adaptive evolution in the Mediterranean isles. *Herpetological J.* 1: 465-473.
- Eisentraut, M. (1930). Beitrag zur Eidenchsenfauna der Pityusen und Columbreten. *Mitt. Zool. Mus. Berlin*, 16: 397-410.
- Klemmer, K. (1961). Las Islas Columbretes - die Schlangeninseln ohne Schlangen. *Natur und Volk*, 91: 39-47.

- Kuranova, V. N. 1989. On melanism in the viviparous lizard and common adder. *Vestn Zool.* 2: 59-61.
- Lovich, J. E., McCoy, C. J. & Garska, W. R. 1990. The development and significance of melanism in the slider turtle, pp. 233-253, in: Gibbons, J. W. (ed.). *Life history and ecology of the slider turtle*. Chap. 19. Smithsonian Institution Press, Washington, D.C.
- Malkmus, K. 1976. Ein negrino der Bergeidechse (*Lacerta vivipara*) aus den schladminger Tauern. *Nachr. Naturwiss. Mus. Aschaffenburg.* 84: 11-16.
- Pérez-Mellado, V. 1984. Sobre un ejemplar melánico de *Podarcis hispanica* (Steindacher, 1870). *Doñana, Acta Vertebrata.* 11: 320-321.
- Pérez-Mellado, V. 1989. Estudio ecológico de la lagartija balear *Podarcis lilfordi* (Günther, 1874) en Menorca. *Revista de Menorca.* 80: 455-511.
- Sage, B. L. 1962. Albinism and melanism in birds. *British birds.* 55: 202-225.
- Zuffi, M. 1986. Su *Podarcis muralis maculiventris* (Werner, 1981) melanica in Risaia a Bereguardo (Pavia). *Atti. Soc. ital. Sci. nat. Mus. Civ. Stor. Nat. Milano.* 127: 293-296.