### **NADA Practice Resource**

# WORKING WITH WOMEN ENGAGED IN ALCOHOL AND OTHER DRUG TREATMENT

#### SECOND EDITION

A resource developed by Network of Alcohol and other Drugs Agencies which aims to support the provision of best practice interventions for women accessing alcohol and other drug treatment









© Network of Alcohol and other Drugs Agencies 2016.

This work is copyright. You may download, display, print and/or reproduce this material in unaltered form only (retaining this notice) for your personal, non-commercial use or use within your organisation. Requests for further authorisation should be directed to the CEO, NADA, PO Box 2345, Strawberry Hills NSW 2012.



NADA is supported by funding from the Australian Government Department of Health.

NADA acknowledges the traditional custodians of country throughout NSW and the ACT and their continuing connection to land, culture and community. We pay our respects to elders past, present and future.

**Suggestion citation:** Network of Alcohol and other Drugs Agencies (NADA). NADA Practice Resource: Working with Women Engaged in Alcohol and Other Drug Treatment (2nd ed). Sydney: NADA; 2016.

First edition published and printed in Sydney by NADA in 2015.

This revised edition published and reprinted in Sydney by NADA in 2016. This edition includes an updated section 3.4 Transgender women.

**Disclaimer:** While every reasonable effort has been made to ensure the contents of this resource are correct, NADA does not accept responsibility for the accuracy or completeness of the contents, and is not liable to any person in respect of anything done or omitted in reliance upon the content of this document.

# 2016

#### **NADA Practice Resource**

# WORKING WITH WOMEN ENGAGED IN ALCOHOL AND OTHER DRUG TREATMENT

#### **SECOND EDITION**

The NADA Practice Resource: Working with Women Engaged in Alcohol and Other Drug Treatment, was developed as part of the Network of Alcohol and other Drugs Agencies (NADA) Women's AOD Services Development Program.

## ABOUT THIS RESOURCE

This resource aims to support the provision of best practice interventions for women accessing Alcohol and Other Drug (AOD) treatment and to effect organisational change around becoming gender responsive, family inclusive and trauma informed. While this resource was developed for use by the non government AOD sector in NSW, the information in it is relevant for, and could be used by, organisations that work with women who present with a range of complex needs.

The Women's AOD Services Development Program was funded by the Department of Health (formerly the Department of Health and Ageing) under the Substance Misuse Services Delivery Grant Fund in July 2013 to improve the capacity of NSW non government alcohol and other drug services to meet the needs of substance using women, with and without children who have mental health and other complex health and social issues.

#### **ABOUT NADA**

The Network of Alcohol and other Drugs Agencies (NADA) is the peak organisation for the non government alcohol and other drugs sector in NSW. Our vision is a connected and sustainable sector providing quality evidence based programs to reduce alcohol and drug related harms to NSW communities.

We represent approximately 100 organisational members that provide a broad range of services including health promotion and harm reduction, early intervention, treatment and aftercare programs. Our members comprise of services that are diverse in their structure, philosophy and approach to alcohol and other drugs service delivery. NADA provides a range of programs and services that focus on sector and workforce development, information management, governance and management support, sector representation and advocacy, as well as actively contributing to public health policy.

NADA is governed by a board of directors elected from the NADA membership. We are accredited under the Australian Service Excellence Standards.

Further information about NADA and our programs and services is available on the NADA website at www.nada.org.au.

#### **ACKNOWLEDGEMENTS**

NADA sincerely thanks each of the 51 women from six services across NSW who generously shared their views and experiences of seeking and receiving treatment for their drug and alcohol issues. NADA would also like to recognise the commitment of the Women's Alcohol and Other Drug Services Network to improving outcomes for women and for their vision that both inspired and guided this important work.

NADA would like to acknowledge the contribution of the following individuals and organisations in the development of this resource:

#### Women's Alcohol and Other Drug Services Network

- Detour House Inc.
- Dianella Cottage
- Guthrie House
- Jarrah House, Women's Alcohol and Drug Advisory Centre
- Kamira
- Kathleen York House, Alcohol and Drug Foundation NSW
- Leichhardt Women's Community Health Centre
- Lou's Place
- Phoebe House
- Sydney Women's Counselling Centre
- New Beginnings, We Help Ourselves
- Women and Girls Emergency Centre

#### The NADA Women's AOD Services Development Program Expert Advisory Group

- A/Prof Lucy Burns, National Drug and Alcohol Research Centre (NDARC), University of NSW
- Dr Courtney Breen, (NDARC), University of NSW
- Sunara Fernando, Mental Health and Drug and Alcohol Office (MHDAO), NSW Ministry of Health
- Jane Fischer, National Centre for Education and Training on Addiction (NCETA), Flinders University
- Alice Hanna, Jarrah House, Women's Alcohol and Drug Advisory Centre
- Kristi Hawkins, Australian Government Department of Health
- Catherine Hewett, Kamira
- Dr Latha Nithyanandam, Kathleen York House, Alcohol and Drug Foundation NSW
- Professor Juanita Sherwood, Faculty of Arts and Social Sciences, University of Technology Sydney

#### Other reviewers and contributors:

- ACON
- Clinical Issues Team, Office of the Senior Practitioner, Department of Family and Community Services
- Drug and Alcohol Multicultural Education Centre (DAMEC)
- The Gender Centre

#### **Resource Development**

- Dr Suzie Hudson of Speak Listen Heal Counselling and Consultancy
- LeeJenn Health Consultants

NADA also acknowledges contribution from Teddy Cook, The Peer Advocacy Network for the Sexual Health of Trans Masculinities (PASH.tm) whose feedback resulted in the updated 3.4 Transgender women section.

# **GLOSSARY OF TERMS**

ASSERTIVE FOLLOW UP

Is a process of checking in with clients via SMS, phone call, email or letter as a means of providing support or following up missed appointments.

**AFTERCARE** 

Is the provision of support and/or services that occur after or at the conclusion of a specific AOD treatment program. It may take the form of a support group, leisure activity, supported accommodation or case management.

**COMPLEX NEEDS** 

Refers to the interaction of the multiple areas of need that people accessing AOD treatment may experience, including mental health issues, experiences of trauma, family and domestic violence, experiences with the criminal justice system, physical and cognitive health needs and factors associated with marginalisation.

FAMILY AND DOMESTIC VIOLENCE

A term that incorporates a wide range of abusive behaviours that occur in the context of intimate and family relationships, involving spouses/de-facto partners, ex-partners, children, siblings, parents and/or caregivers.

FAMILY SENSITIVE OR FAMILY INCLUSIVE PRACTICE An approach that recognises that interventions are more effective when they include family members, where the term family refers to all supports or significant others. It supports the view that individuals influence other members in their environment – especially family – and that family members, in turn, have an impact on these individuals.

**GENDER RESPONSIVE** 

Gender responsiveness, in an AOD context, can be defined as aiming to understand and address issues experienced by women and creating an environment that reflects these understandings through approach and treatment options.

HOLISTIC PRACTICE

Is an approach to providing care in a way that considers all aspects of a clients life, including their physical and mental health, living situation and social networks.

POST TRAUMATIC STRESS DISORDER

Post-traumatic stress disorder is a specific set of responses that can develop in people who have been through a traumatic event. The traumatic event(s) experienced or witnessed by an individual may have threatened their life or safety, or that of others around them, and led to feelings of intense fear, helplessness or horror. This can be a serious accident, physical or sexual assault, war or torture, or disasters such as bushfires or floods.

STRENGTHS BASED APPROACH

Recognises the resilience of individuals and focuses on their strengths and potential rather than their deficits.

TRAUMA INFORMED CARE/PRACTICE

Term used to describe a way of working with people that acknowledges the lasting impact of trauma. Trauma informed care also emphasises physical, psychological and emotional safety for both clients and providers, and helps survivors regain a sense of control and empowerment over their lives.

# **ACRONYMS**

AIHW	Australian Institute of Health and Welfare	
AOD	Alcohol and Other Drug(s)	
BBV	Blood Borne Virus	
CALD	Culturally and Linguistically Diverse	
СВТ	Cognitive Behavioural Therapy	
CRA(FT)	Community Reinforcement Approaches (and Family Training)	
СТО	Community Treatment Order	
DAMEC	Drug and Alcohol Multicultural Education Centre	
FASD	Fetal Alcohol Spectrum Disorders	
FDV	Family and Domestic Violence	
FACS	Family and Community Services	
LGBTI	Lesbian, Gay, Bisexual, Transgender, Intersex	
MHDAO	Mental Health and Drug and Alcohol Office	
NADA	Network of Alcohol and other Drugs Agencies	
NCETA	National Centre for Education and Training on Addiction	
NDARC	National Drug and Alcohol Research Centre	
NDSHS	National Drug Strategy Household Survey	
NGO	Non government organisation	
NIDAC	National Indigenous Drug and Alcohol Committee *Defunded in 2014	
ООНС	Out of Home Care	
PTSD	Post Traumatic Stress Disorder	
STI	Sexually Transmitted Infections	

# CONTENTS

How to navigate this resource	vii
Purpose of the resource	vii
SECTION 1: WOMEN IN ALCOHOL AND OTHER DRUG TREATMENT	1
1.1 Women and their experience of alcohol and other drug use	
1.2 Unpacking issues and identifying needs of women in alcohol and other drug treatment	
1.3 Considerations for women on pharmacotherapy	
1.4 Women accessing treatment for a significant other's drug use	
SECTION 2: CREATING AN ENABLING ENVIRONMENT FOR WOMEN: BEST PRACTICE APPROACHES	
FOR AOD TREATMENT	
2.1 Establishing a gender-responsive service	
2.2 Adopting a strengths-based approach to treatment provision	19
2.3 Trauma informed care and trauma specific approaches	20
2.4 Family-inclusive practice	23
2.5 Evidence based interventions and coordinated case management	
2.6 Women's sexual health and safety in AOD treatment	28
2.7 Assertive follow-up, outreach and aftercare	30
SECTION 3: IMPROVING ACCESS AND RETENTION FOR WOMEN FROM DIVERSE BACKGROUNDS	31
3.1 Working effectively with Aboriginal and Torres Strait Islander women	31
3.2 Increasing supports for culturally and linguistically diverse women	35
3.3 Enhanced service-provision for women from LGBTI communities	38
3.4 Transgender women	40
SECTION 4: PREGNANCY AND PARENTING IN AOD TREATMENT	43
<b>4.1</b> Alcohol and other drug use in pregnancy: Improving pre-natal care outcomes	44
<b>4.2</b> Introducing child-sensitive service provision and support for positive parenting	47
4.3 Working effectively with Family and Community Services	49
SECTION 5: BUILDING STRONG PARTNERSHIPS TO INCREASE POSITIVE OUTCOMES FOR WOMEN	55
5.1 Involving women in effective case management	55
<b>5.2</b> Building community-based partnerships	57
5.3 Identifying partnerships that support effective aftercare	59
SECTION 6: ORGANISATIONAL CHANGE AND SELF-CARE FOR WORKERS	61
6.1 Areas for review and enhancement in mixed-gender organisations	63
<b>6.2</b> Enhancing organisational approaches that promote self-care	66
SECTION 7: RESOURCES AND GUIDELINES	69
7.1 Information and support	69
7.2 Practice resources and toolkits	74
7.3 Government guidelines and strategies	79
7.4 Training and workforce development centres	81
SECTION 8: REFERENCES	83

#### Do you work in the AOD sector?

Chances are you bring some essential qualities to your work, including:

- a non-judgemental attitude
- empathy and compassion
- a willingness to work with clients 'where they are at'
- the ability to listen and support clients and their loved ones
- a commitment to challenge discrimination and stigmatising language and behaviour
- skills and professionalism
- an optimistic approach that gives your clients hope
- a holistic practice that considers all of the factors that contribute to improved wellbeing
- a passion for social justice and equality
- a curiosity that drives you to engage with innovative, best practice interventions.

The great news is that these are exactly the kinds of qualities that are essential when working with women who experience alcohol and other drug use issues.

We know that by the time women access (AOD) treatment they have faced multiple challenges in their lives. However, the multitude of complex needs with which women may present must never overshadow the strength it has taken to make that first step, and the resilience they have demonstrated in getting this far.

If you do nothing else in the course of your important work alongside these courageous women, simply acknowledge the achievement they have made in finding their way to you.

This resource is intended to support you in the work that you and your organisation engage in every day. In this resource you will find:

- snapshots of specific challenges that women face in the context of treatment
- frontline worker experiences from organisations that provide specialist AOD services for women
- case studies and best practice approaches that you and your organisation can implement.

While this resource covers a wide range of topic areas, it is by no means an exhaustive guide – important original documents have been consulted and made reference to within the resource and you are encouraged to seek them out for more comprehensive information.

# HOW TO NAVIGATE THIS RESOURCE

This resource is designed to be a quick reference guide. It is separated into eight sections that cover a range of specific topic areas. The concluding section provides important additional guidelines and recommendations for further reading, and is grouped according to the topic areas they relate to.

Sections have been structured in order to promote thought, reflection and discussion around practice approaches to working with women in the AOD treatment sector. Each section addresses a key area of knowledge that contributes to improved ways of working. Concise summaries of research and practice literature relevant to each topic area are presented throughout the resource and in addition, a number of regular features are used including:



**Key points and principles** – appear at the beginning of each section for quick reference.



**From the frontline** – provides a snapshot of real-life experiences of women in specialist women's services and shares and practice wisdom sourced from workers based in either women-specific AOD services or mixed gender services who were consulted as part of the development of this resource.



**Case studies** – present lived experiences of women accessing AOD treatment and provides examples of the complex array of issues and challenges faced by women within the context of a treatment journey.



**Find out more** – provides references and links to original source literature, published research, guidelines and clinical guides that have been reviewed in the development of this resource. Recommended resources can be used as a catalyst for further reading and exploration into material that will support improved treatment provision and outcomes for women and their loved ones. In **Section 7** and **8** of this resource are specific toolkits, guidelines and references for finding out more about all the material that has informed the resource.

# PURPOSE OF THE RESOURCE

The purpose of the resource is to raise the profile of women in AOD treatment. Improving service provision to women means acknowledging their unique experiences and perspectives, enhancing the best practice approaches AOD treatment services already have in place and/or adopting philosophical perspectives that give a greater voice to the needs of women. Organisation level engagement with interventions that serve the needs of women have far-reaching benefits. These include improving gender responsiveness, addressing issues related to family violence for both victims and perpetrators, raising awareness regarding the needs of children and introducing concepts around positive parenting and family inclusive practice.

This resource provides structural solutions, such as building sustainable community partnerships, professional development and worker self-care that are essential in supporting improved work practices with women. Developed from evidence-based practice and research, this resource has brought together relevant findings from the literature and practice experience as a starting point for deeper consideration and discussion into the needs of women accessing AOD treatment.



# 1

# WOMEN IN ALCOHOL AND OTHER DRUG TREATMENT

# 1.1 Women and their experience of alcohol and other drug use



- Gender shapes substance use and misuse in relation to initiation, access, consumption, experiences of dependence, AOD-related risk behaviours and AOD treatment outcomes.
- Women access AOD treatment at lower rates than men and are under-represented in the drug and alcohol treatment system (according to Australian and international data).
- A range of issues create barriers for women's access to AOD services, including social stigma, discrimination, experiences of trauma, childcare and child custody concerns, and financial issues.
- On entering AOD treatment, women present with higher rates of: mental health issues; experience of complex trauma as a result of childhood physical and sexual abuse and/or family and domestic violence; AOD-related risk taking; pregnancy and childcare issues; and greater social and economic disadvantage.
- Pregnancy and parenting are specific areas of need that require effective support and intervention for women engaged in problematic substance use. The key to engaging women is to reassure them that there is 'no wrong door'.
- Pharmacotherapy for women works best when implemented with other supportive interventions such as coordinated case management, counselling, group therapy and practical support.
- A high proportion of women who access AOD treatment are seeking assistance for the problematic substance use by a loved one – principles of family inclusive practice are best practice in this context.

WOMEN'S AOD USE AND ENGAGEMENT WITH TREATMENT IN AUSTRALIA Women's substance use and their experience of problematic use and dependence are shaped by gender. Australian social and environmental influences also have a significant role to play in terms of accessibility, consumption norms and the social perceptions that are attached to different drug types. However gender is the dominant lens through which substance use and misuse is judged. Building recognition of the ways in which gender shapes women's experiences of substance use and treatment for problematic use has the potential to improve ways of working in the AOD sector, and the treatment outcome of improved general wellbeing for women in Australian society.

AOD use is generally more prevalent among males than females, and in Australia overall, females are far less likely than males to use either licit or illicit substances (AIHW 2014). Gender differences are evident in most patterns of substance use and females in the National Drug Strategy Household Survey (NDSHS) 2013 were found to be around half as likely as males to either drink daily, or to have injected any drug in the past year. The main exception to this pattern is the use of pharmaceuticals and prescription drugs for non-medical purposes. An equal number of men and women reported having used pharmaceuticals such as analgesics, sedatives, steroids or opiates in a non-prescribed way in the past 12 months (AIHW 2014).

WOMEN'S AOD USE AND ENGAGEMENT WITH TREATMENT IN AUSTRALIA

continued

In Australia, the estimated prevalence of harmful substance use or dependence disorders among women is around half that of men. However, in 2012/13, monitoring of AOD treatment utilisation showed that less than one third of treatment episodes were for female clients seeking help for their own substance use (AIHW 2014). Alcohol is the most common principal drug of concern for both women and men seeking treatment, and male clients made up the majority of treatment episodes in 2012/13 where alcohol, cannabis, heroin and amphetamine were identified as the primary drug. In contrast, almost half of the episodes related to benzodiazepines as the primary drug were with women (AIHW 2014).

Consistent with national rates that suggest women are unrepresented in treatment, 33% of NSW treatment episodes at over 250 AOD services were for female clients, whose median age was 40 years (AlHW 2014). Women appear to be seeking and entering treatment for their substance use at lower rates than men, when they are older and at rates that are disproportionate to their likely level of need. Subsequently, identifying and responding to women's treatment needs and women-specific barriers to accessing services may increase rates of treatment utilisation.

Women with problematic substance use and those who enter treatment have been found to differ from male substance users in a range of ways. A number of factors and characteristics have been found to be associated with their increased risk of initiation to substance use, increased risk of transition to problematic substance use and dependence, as well as their outcomes once engaged in treatment. Risk factors that are highly prevalent and, in some cases, unique to women substance users need to be taken into account in order to deliver best practice treatment. The literature also indicates that women accessing AOD treatment are more commonly poorly educated and have low levels of income, have a financial dependence on partners or others and have lower rates of employment than their male counterparts (McCabe et al. 2013, Yeom 2011).

Gender differences are evident among injecting drug users, where women who inject drugs are more likely than men to inject with multiple partners, share paraphernalia after an injection partner, exchange sex for money or drugs, and have difficulty negotiating condom use with their sex partners (Strathdee et al. 2001; Ashley et al. 2003) resulting in vulnerability to HIV infection.

An awareness of the challenges that women face in engaging in treatment for their problematic use of alcohol and other drugs provides the foundation from which to develop appropriate interventions and make improvements to service provision. Explored in the next section are some of the areas of complexity that women may experience on their journey into treatment.

# 1.2 Unpacking issues and identifying needs of women in AOD treatment

Improving the access and retention of women in AOD treatment can be facilitated by an awareness of the factors that may have contributed to their problematic alcohol and drug use, and those that perpetuate it. Women generally present with a more severe clinical profile than men when entering treatment, with more problems related to mental health, family and interpersonal relationships, employment, and physical health (Green 2006), including involvement with child protective services and partners who are also substance users (Brady & Ashley 2005). In spite of the fact that women who enter treatment generally have fewer years of substance use and have used smaller quantities than men, they present with an equivalent severity of substance use problems and symptoms (Covington et al. 2008, Yeom 2011).



Many of the issues and needs of women accessing AOD treatment identified through research are echoed by the AOD sector. According to key informants from NSW non government AOD treatment services who were consulted in the development of this resource, areas of specific need include:

- pharmacotherapy the lack of places for women on pharmacotherapy programs and inconsistent collaborative care planning with prescribers and pharmacotherapy case managers
- parenting issues a lack of treatment places for women with children in their care; increasing numbers of women presenting for treatment with children in out-of-home care (OOHC); inconsistent collaborative care planning with case workers from Family and Community Services (FaCS)
- increasing complexity of presentations trauma and its enduring effects on AOD use and women's quality of life, mental health issues, cognitive impairment that complicates drug and alcohol treatment, physical health issues particular to women
- housing issues particularly the lack of affordable and appropriate housing available to women
- changing cohorts of women seeking treatment, such as greater numbers of first-time treatment seekers and well-resourced older women.

## STIGMA AND DISCRIMINATION

Several theories relate to the stigma and discrimination experienced by women who use alcohol and other drugs, and they usually align with traditional views of women's roles in society. According to Shimmin "stigma for women who use any licit or illicit drugs is more severe than for men because of women's "place" in society, as those who bear and rear children and who are seen to uphold the moral and spiritual values of society. There is also the negative stereotype that women users are sexually promiscuous because of their drug or alcohol use. This association is not seen in men." (2009).

"Stigma for women who use any licit or illicit drugs is more severe than for men because of women's "place" in society..." Shimmin, 2009

Women often face social barriers as they are more likely than men to encounter disapproval for their substance use and their participation in treatment. This may include the loss of family support due to substance use or associated criminal behaviours (Ashley et al. 2003, Taylor 2010).

While stigma is potentially of concern for all substance users, women with drug and alcohol use issues face a significant social stigma and discrimination in relation to pregnancy and parenting. Along with the barriers posed by family responsibilities or lack of childcare options if they were to seek treatment, the stigma of being a substance-using mother and the fear of losing custody of children are significant barriers to treatment entry and engagement (Greenfield et al. 2007).

# -rom the frontline



Find out more...about tackling stigma and discrimination.

#### NUAA's Stigma and Discrimination Campaign

In order to address issues and challenge issues of stigma and discrimination, NUAA has been implementing training based on resources developed by the Australian Injecting & Illicit Drug Users League (AIVL). NUAA targets health care professionals and those working in the drug and alcohol area as well as the broader community and people who inject drugs themselves to tackle stigma and discrimination in a thought-provoking and innovative manner.

Website: www.nuaa.org.au www.aivil.org.au

#### Developing a Stigma Reduction Initiative SAMHSA 2006 (Pub No. SMA-4176)

This resource kit is intended to raise awareness of mental health and help counter the stigma and discrimination faced by people with mental illnesses. It is designed to support the activities of those who plan to mount a statewide, regional, or local effort to address and counter stigma and discrimination. It is intended for use by local mental health advocates, consumers of mental health services and their family members, community leaders, and other organisations and individuals who have dedicated themselves to eliminating the barriers of stigma and discrimination. It contains resources and templates.

#### Stigma Watch SANE Australia

SANE StigmaWatch reflects and acts on community concern about media stories, advertisements and representations that stigmatise people with mental illness or inadvertently promote self-harm or suicide.

Website: www.sane.org/stigmawatch

#### Tackling the Stigma: Drugs and Discrimination. Of Substance July 2012 (10:12)

Of Substance is an Australian National Council on Drugs (ANCD) initiative and is produced with funding from the Australian Government Department of Health. Written in plain English, Of Substance reports on the latest news and research in the field; regional, national and international data and trends; and a range of skills and techniques for best practice treatment. This particular issue focuses on stigma experiences of those who use alcohol and other drugs.

CO-OCCURRING MENTAL HEALTH AND AOD ISSUES In Australia, 20-30% of women who have alcohol use issues meet diagnostic criteria for a mental health disorder (ABS 2008). Women who use alcohol and other drugs have a higher prevalence of co-occurring mental health issues such as mood, anxiety, post-traumatic stress disorders (PTSD) and eating disorders than men (Covington et al. 2008). Affective disorders, such as major depression, are associated with poorer treatment outcomes and higher rates of relapse. In a review of the evidence for women in drug and alcohol treatment, Greenfield et al. (2007) suggest that higher rates of co-occurring disorders among women, such as depression, anxiety, eating disorders and PTSD, may make it difficult for women to gain appropriate treatment for both mental health and substance use problems. Additionally, the experience of multiple disorders may increase the likelihood that women perceive their issues as specific to their mental illness with the result that they focus on seeking psychiatric help, rather than AOD treatment.

In a study of over 300 women experiencing co-occurring substance use and PTSD, those who perceived their need for psychological and emotional treatment as highly important were found to be significantly more likely to not attend, or drop out early from, AOD-related treatment (Resko & Mendosa 2013). Among women substance users, common co-occurring issues such as depression may go undetected, and women may also choose to self-medicate rather than seek professional assistance.

The complexities of some highly prevalent mental health issues present serious additional risks. Alcohol and drug use is strongly associated with eating disorders for women, and substance use and eating disorders are both associated with the highest mortality risks of all psychological disorders. Women who are nutritionally deprived when they present to AOD treatment are best supported with medical intervention and advice because individuals with substance dependence and an eating disorder face increased problems with withdrawal and dependency (Harrop & Marlatt 2010).



AOD treatment services who were consulted in the development of this resource reported that they were witness to an increasing complexity of mental health problems experienced by the women who presented to their services. When asked to nominate the main mental health issues that affect female clients of their service, the following issues were nominated: depression; anxiety; trauma; personality spectrum disorders; psychotic disorders and eating disorders.

All of our women have mental health issues – a lot are on antidepressants. All of them are depressed and anxious. Some have PTSD and the majority of women have had a major trauma in their lives. Recently, we had a woman referred who had just come out of hospital for drug-induced psychosis.

We ask for a mental health report before the client comes in but we sometimes find in the withdrawal period that the alcohol and drug use has been protective of the mental health problem so we will see psychosis emerging.

We have a lot of women on mood stabilisers and we have had a couple of women who've come to us on CTOs (Community Treatment Orders)... and went very well actually. We had one young woman who stayed... for five months... but we've had other women admitted with psychosis and they struggled to live in that sort of [residential treatment] community setting.

Due to the high risks and complex needs of women experiencing substance use disorders and co-occurring mental health issues, integrated and holistic treatment approaches are recommended. There's also a need for a longer-term approach to recovery and sensitive outreach, engagement and support interventions (Brown & Melchior 2008, Chen & Biswas 2012). See **Section 2.7** for more information on outreach as an intervention.



**Find out more...**about working with co-existing mental health and alcohol and other drug issues.

Guidelines on the management of co-occurring alcohol and other drug and mental health conditions in alcohol and other drug treatment settings. NDARC 2010 (Mills et al.)

The purpose of these guidelines is to provide AOD workers with up-to-date, evidence-based information on the management of comorbid mental health conditions in AOD treatment settings. They are based on the best available evidence and draw upon the experience and knowledge of clinicians, researchers, consumers and carers.

Complex Needs Capable: A Practice Resource for Drug and Alcohol Services NADA 2013

Complex Needs Capable and the accompanying website was developed to build capacity within the non government drug and alcohol sector by supporting the development of staff skills, knowledge and confidence, and of organisational capacity in responding to the needs of people with drug and alcohol issues and complex needs such as cognitive impairment and criminal justice system contact.

Website: www.complexneedscapable.org.au

# NSW Clinical Guidelines: For the care of persons with comorbid Mental Illness and Substance Use Disorders in Acute Care Settings NSW. NSW Health 2009

These guidelines have been written for those working in the AOD and/or mental health sectors who provide care for people with comorbid mental health and substance use disorders.

## RELATIONSHIP ISSUES

Personal relationships can affect women's AOD treatment entry and success, representing treatment seeking barriers (Greenfield et al. 2007, Grella 2009, Tuchman 2010). Problematic intimate relationships have the potential to increase the risk and harms women experience as a result of their substance use. On the other hand, more stable or supportive relationships can reduce these risks and harms. Substance misuse by significant others, and living with a substance user, have been found to be risk factors associated with poorer outcomes and higher likelihood of relapse for women in treatment, but this relationship has not been demonstrated as significant for men (Grella et al. 2008).

Personal relationships and family history appear to be a more significant influence on women's initiation, pattern of use, and continuation of problematic alcohol and drug use than for men. For example, women with alcohol and drug problems appear more likely than men to have been raised in a family environment of heavy drinking or problematic drug use (Ashley et al. 2003, CSAT 2009). Furthermore, women who identify as substance users are more likely to have intimate partners who are substance users (Brady & Ashley 2005) and appear more likely to report that they had family, friends or partners who used drugs or who supported their continued substance use (Greenfield et al. 2007).

Some women perceive their entry into drug use, or their continued use with their partner as a way to maintain or develop more intimate relationships (Kay et al. 2010). Furthermore, situations and conditions such as having a stressful marriage and being pressured to use drugs by sexual partners have been found to significantly affect women's post-treatment relapse to alcohol use (Grella 2008).

The principles of family inclusive practice (explored in more detail in **Section 2.4**) are applicable when considering engaging partners of women who experience substance use issues. While in some cases it may not be appropriate to involve partners directly, referrals to alternative supports such as family groups or counselling is recommended. Education and information about the process of dependent substance use, the related issues and what is involved in treatment can assist those supporting women accessing AOD treatment. Confidentiality is key in working with significant others. However, clear explanations about what confidentiality means in a treatment context at the first point of contact is best practice, while being empathetic to the reasons for requesting information.



Find out more...about strengthening supportive intimate relationships.

#### Relationships Australia

Relationships Australia is a leading provider of relationship support services for individuals, families and communities. Services are for all members of the community, regardless of religious belief, age, gender, sexual orientation, lifestyle choice, cultural background or economic circumstances.

Website: www.relationships.org.au

#### Family Drug Support

Family Drug Support (FDS) was formed in 1997 after its founder, Tony Trimingham lost his son to a heroin overdose. FDS is a caring, non-religious and non-judgemental organisation, primarily made up of volunteers who have experienced firsthand the trauma and chaos of having family members with drug dependency. FDS provides a 24 hour/7 day a week telephone support line, website, support groups and courses for family members and significant others.

Website: www.fds.org.au

# **Families and friends affected by the drug and alcohol use of someone close.** NSW Health 2004

Answers common questions and concerns of those who are experiencing the challenges of having someone they care about engage in problematic substance use. These resources are also available in Arabic, Chinese, Turkish, Serbian, Croatian and Vietnamese from the multicultural health communications website.

Website: www.mhcs.health.nsw.gov.au

#### FAMILY AND DOMESTIC VIOLENCE (FDV)

Women are more likely than men to identify relationship issues as a cause for their problematic substance use. In addition, women in treatment have much higher rates of family violence experience than women in the general community (Greenfield 2010). A high prevalence of intimate partner violence among substance using couples suggests that where drug use is central to a couple's relationship, women's ability to enter treatment may also be limited by the threat of violence. Family and domestic violence encapsulates a wide range of abusive behaviours, occurring within intimate and familial contexts. While FDV predominantly involves male violence against their female partners, other relationship violence is included in this definition such as that existing in same-sex relationships, non-spousal relationships and carer relationships – parent to child and vice versa (White et al. 2013).

"Families would benefit from AOD services making the links between AOD and FDV explicit. This has the potential to increase the client's capacity and motivation to change AOD use, with improved outcomes for their children." (White et al. 2013).

Given that a considerable proportion of people accessing alcohol and other drug treatment are men, there is potential for some of them to have been perpetrators of FDV. This presents an opportunity in the context of AOD treatment to explore FDV as standard practice, and an issue that is not just targeting women who have been victims. One of the key recommendations from Australia's national research centre on AOD workforce development (National Centre for Education and Training on Addiction), resource "Can I Ask? – An alcohol and other drug clinician's guide to addressing family and domestic violence" (White et al. 2013) is to include training that builds capacity in the area of FDV so that AOD workers feel confident to raise FDV issues, to provide screening and assessment and explore risk assessments and safety planning.



Find out more...about addressing FDV in AOD settings.

# Can I Ask? An alcohol and other drug clinician's guide to addressing family and domestic violence. NCETA 2013 (White et al.)

This resource explores the relationship between AOD and FDV, with a focus on identifying how the AOD sector can better support clients who have co-occurring AOD and FDV issues, and minimise associated harms experienced by their children.

# Breaking the silence: Addressing family and domestic violence problems in alcohol and other drug treatment practice in Australia. NCETA 2012 (Nicholas et al.)

This review explores the relationship between AOD and FDV services, with a focus on identifying how the AOD sector can better support their clients who have co-occurring family and domestic violence issues and minimising the harm experienced by children.

**The Domestic Violence Line** provides telephone counselling, information and referrals for women and same-sex partners who are experiencing or have experienced domestic violence. Trained female caseworkers are sensitive to the needs of people who have experienced domestic violence. Domestic Violence Line staff are aware of the needs of Aboriginal women and women from other cultures, as well as those living in rural and remote areas. Interpreters and TTY can be arranged where necessary to ensure that all people, regardless of their language or disability can use the service. The service has an extensive list of contacts, people and services across NSW who can help.

Phone: 1800 656 463

Website: www.domesticviolence.nsw.gov.au

# FINANCIAL DISADVANTAGE AND HOMELESSNESS

Women entering treatment commonly experience lower levels of education and income, and have lower rates of employment. Women have more problems related to lack of employment and vocational skills (Grella 2008) and report financial dependence on partners or others (Engstram et al. 2012). Furthermore, low socio-economic status and income inequality are often linked with poor health and wellbeing and have been associated with serious drug-related harms including fetal alcohol syndrome, alcohol-related deaths and drug overdoses (Spooner & Hetherington 2004).

Homeless women who use alcohol and drugs have a higher risk of mental health issues, experience of trauma and potential involvement in the sex industry (Brown et al. 2012, Hopper et al. 2010). They are vulnerable to further poverty, family breakdown and violence (Dawson et al. 2013). Addressing these issues, in addition to providing material support, is therefore highly important for this group.

Positive socio-economic factors, such as being gainfully employed, having a higher income and higher level of education generally mean better rates of treatment retention (Greenfield et al. 2007). Engagement in AOD treatment is an opportunity to explore further education and vocational training options with women and reaffirms the benefit of partnership building with organisations that specialise in this area. See **Section 5** in this resource for more information.



From the frontline (

Homelessness or the risk of homelessness is of great concern to women accessing AOD treatment, particularly given that prolonged problematic substance use is a significant contributing factor to financial difficulties and accommodation instability. Many AOD services have identified inconsistent or poor access to support services needed by women such as safe and stable housing.

Homelessness is on the top of the list [of concerns]. When women lose their accommodation – they end up with someone just to get a roof over their head. They sleep on couches and they trade the need for safety.

Housing is a massive gap. It's difficult for women with more than one child, single women... if they are on pensions their ability to pay commercial rent is difficult.

There is a population of women who want to get out of the criminal justice system and they are stuck and unable to get out. The women have been travelling [...] from the city to get away from their old networks but there's few supports here, there's no housing.

Providing assertive community outreach has helped identify people who are homeless and educate them about available services and AOD treatment (CSAT 2013). The literature suggests that targeted informal day programs and drop-in programs for homeless women with substance use issues are effective. Particularly where they are successfully integrated components of harm reduction, trauma informed services and social support to homeless and marginally housed. Studies indicate that low-threshold and tailored environments for care can engage highly vulnerable and intervention resistant clients into regular service contact, with benefits for health and wellbeing (Hopper et al. 2010, Magee & Huriaux 2008). Read more about the effectiveness of outreach in **Section 2.7** of this resource.



**Find out more...**about responding to homelessness in AOD settings.

**Specialist Homelessness Services Case Management Resource Kit** NSW Dept. of Family and Community Services 2012

This resource kit is designed to give SHS services up-to-date guidance on the delivery of case management that reflects evidence-based best practice and aligns with new Commonwealth and state policy directions in responding to homelessness. The kit aims to provide caseworkers and casework supervisors across all SHS with knowledge and tools to work effectively with people who are homeless or at risk of homelessness.

Trauma-Informed Organizational Toolkit for Homelessness Services (Guarino et al. 2009)

This Toolkit offers homeless service providers with concrete guidelines for how to modify their practices and policies to ensure that they are responding appropriately to the needs of families who have experienced traumatic stress.

# CRIMINAL JUSTICE ISSUES

Women involved in the criminal justice system are understood to generally experience high levels of social disadvantage. Among police detainees in Australia, female offenders were found to be substantially more socially and occupationally disadvantaged than women in the general community, and more disadvantaged than male detainees (Loxley & Adams 2009). Male detainees were more than twice as likely as women to report having recent income from paid employment, while females were significantly more likely to be receiving government welfare and to have dependent children in their home (Loxley & Adams 2009). Women involved in the criminal justice system experience high rates of victimisation and mental health issues (Tripodi et al. 2011).

Hall et al. (2013) recommended that interventions should comprehensively address the psychological and social needs of women involved in the criminal justice system, because addressing victimisation, drug and alcohol use, and other mental health disorders is more effective than programs that focus on criminal recidivism alone. Best practice interventions for women who have had experience with the criminal justice system should include components that address women's immediate needs, such as assistance in accessing safe and affordable housing and providing job training, education and employment assistance.

Providing an environment that is sensitive to the experiences of institutionalisation by providing space, time and emotional safety will facilitate engagement in AOD-specific treatment interventions (NADA 2013). For drug and alcohol treatment among women in residential correctional facilities, the therapeutic effects of treatment programs appear to be enhanced when trust-based relationships are established, care is individualised and the treatment facilities are separate from the general prison environment (Fingeld-Connett & Johnson 2011).

Women leaving prison face particular challenges and it is understood that on release, women with substance use issues need both practical and emotional support to re-enter the community. This population have significant and complex health care needs, but this time of transition is indicated to be an effective opportunity to offer support, services, and other health-promoting interventions (Colbert et al. 2013). It is useful to note that many of the factors associated with relapse into alcohol and drug use mirror those associated with re-offending.



**Find out more...**about responding to women who have experience with the criminal justice system.

#### Complex Needs Capable: A Practice Resource for Drug and Alcohol Services NADA 2013

Complex Needs Capable and the accompanying website was developed to build capacity within the non-government drug and alcohol sector by supporting the development of staff skills, knowledge and confidence, and of organisational capacity in responding to the needs of people with drug and alcohol issues and complex needs such as cognitive impairment and criminal justice system contact.

Website: www.complexneedscapable.org.au

**Your Guide to Surviving on the Outside** Community Restorative Centre NSW 2007 This booklet is a guide for ex-prisoners and prisoners who are about to be released.

Website: www.crcnsw.org.au

#### WIPAN (Women In Prison Advocacy Network)

WIPAN helps women who have left prison or are close to being released adjust back into the community.

Website: www.wipan.net.au

#### Sisters Inside

Sisters Inside is an independent community organisation, which exists to advocate for the human rights of women in the criminal justice system, and to address gaps in the services available to them.

Website: www.sistersinside.com.au

### COGNITIVE IMPAIRMENT

The term cognitive impairment is used to describe deficits in cognitive skills or functioning, and refers to symptoms present at birth and or acquired later. Deficits related to Fetal Alcohol Spectrum Disorders (FASD), acquired brain injury, and genetic abnormalities are often grouped under this term. The Complex Needs Capable resource developed by NADA explores in detail issues related to cognitive impairment (specifically, intellectual disability, acquired brain injury and FASD), and, more importantly, the strategies and interventions required for services to become more 'capable' in responding to these needs.



The AOD treatment services consulted in the development of this resource highlighted cognitive impairment as a significant issue that affected women clients accessing their service. They suggested that cognitive impairment, though common, is often missed in assessment prior to accessing AOD treatment.

There is a big issue with cognitive impairment in drug treatment settings and people are discharged for non-compliance when in fact they may not be able to understand [what has been asked of them].

Having a cognitive impairment can result in a person not being able to access or maintain the drug and alcohol treatment they require. People with a cognitive impairment often:

- don't receive the services they need
- are unable to maintain waiting list protocols
- fall through services gaps and experience a 'referral merry-go-round'
- are screened out of entry to services
- are perceived to have no or low motivation
- are put in the 'too hard basket'
- are said to have 'challenging/problem behaviours' when they, in fact, have a cognitive impairment
- are exited from services early
- are unable to access services because they don't have a diagnosis
- are told that disability services should address their drug and alcohol issues.

When armed with knowledge about how cognitive impairment can affect a person and the practical strategies and adjustments in service provision, AOD treatment services can successfully support people with intellectual disability, acquired brain injury and FASD to receive the expert drug and alcohol treatment they need (NADA 2013).



**Find out more...**about working with women who have cognitive impairment.

#### Complex Needs Capable: A Practice Resource for Drug and Alcohol Services NADA 2013

Complex Needs Capable and the accompanying website was developed to build capacity within the non-government drug and alcohol sector by supporting the development of staff skills, knowledge and confidence, and of organisational capacity in responding to the needs of people with drug and alcohol issues and complex needs such as cognitive impairment and criminal justice system contact.

Website: www.complexneedscapable.org.au

#### Looking Forward: Acquired Brain Injury. arbias 2011

Looking forward is a handbook for supporting people with acquired brain injury. It gives a basic introduction to brain injury and the challenges involved for people living with acquired brain injury. It offers practical advice and strategies to enable practitioners to empower people to achieve their full potential.

Website: www.arbias.org.au

## PREGNANCY AND PARENTING

Women accessing treatment may be pregnant or become pregnant, presenting unique perinatal risks that are associated with a higher rate of obstetric complications and poor outcomes, including miscarriage, birth defects, decreased birth weight, cognitive deficits, growth restriction, increased neonatal mortality and maternal health problems (Keegan et al. 2010). The negative impacts of substance use on the development of a child in utero and growing up in households where problematic alcohol and drug use are commonplace, are well documented.

However, the key message should be 'no wrong door'. If a woman presents to an AOD treatment service that may not have the capacity to respond to all the issues relating to pregnancy and parenting, then all effort should be made to engage with her while additional or specialist services can be sought. Access to childcare, prenatal care, mental health services, women-only programs, psycho-education sessions focused on women-specific topics and comprehensive services that offer multiple components have all been associated with improved outcomes for women. Pregnancy and care of children can be a significant motivating factor for AOD treatment engagement and retention. However, child safety should always be paramount. Pregnancy and parenting including child safety concerns are explored in more detail in **Section 4** of this resource.



Find out more...about working with women who become/are pregnant.

# Clinical Guidelines for the Management of Substance Use During Pregnancy, Birth and the Postnatal Period NSW Ministry of Health 2014

These guidelines support and provide best practice advice to health professionals in the management of drug use during pregnancy, birth and early development years of the newborn.

# "It's time to have the conversation": Understanding the Treatment Needs of Women Who Are Pregnant and Alcohol Dependent FARE 2013 (Burns & Breen)

This report puts forward the gold standard for the treatment of alcohol dependence in pregnancy. This research reports on information gathered from 11 semi-structured qualitative interviews with clinicians that treat pregnant women with problematic alcohol use provided expert clinical advice on current treatment practices and the gold standard for treatment and factors that impinge on this approach.

# Responding to the Needs of Children and Parents in Families Experiencing Alcohol and Other Drug Problems ADF 2012 (Gruenert & Tsantefski)

This issues paper provides a summary of current research as well as professional opinion relevant to working with families, especially when a parent presents for treatment or support. The paper aims to be a useful resource for both service providers and practitioners, providing tips and advice on family sensitive and family inclusive practice, with a specific focus on supporting children.

#### 1.3 Considerations for women on pharmacotherapy

Women in pharmacotherapy treatment have the same needs as women in drug and alcohol treatment more generally, including a strengths-based approach that is trauma informed, access to childcare and greater focus on engagement and retention in treatment. Opioid dependence is associated with a range of physical health problems and frequent incidence of comorbid mental health issues, and these issues commonly appear to be more prevalent and/or more severe for women (Kay et al. 2010). Compared to male opioid users, female users and women in treatment are significantly more likely to have a mood or anxiety disorder (Grella et al. 2009) and have been found to have more extreme problems related to recent employment, medical, psychiatric, and family social matters (Kelly et el. 2009).

Women participating in methadone programs often have experience of childhood sexual abuse, childhood physical abuse, lifetime experience of intimate partner violence and low levels of financial independence (Engstrom et al. 2012). Opioid users engaged in treatment have demonstrated large improvements in their health and social functioning, but these improvements have been shown to be moderated by the experience of co-existing psychiatric disorders (Grella 2009), highlighting the need for treatment to identify and address women's highly prevalent mental health issues in conjunction with pharmacotherapy.

In situations where women accessing treatment are pregnant, induction to methadone maintenance as part of comprehensive treatment programs is considered best practice (Keegan 2010, Minozzi et al. 2008). Commencement of methadone should ideally take place in an outpatient setting, with maternal and obstetric supervision - as is common practice in Australia, and supported by the American College of Obstetricians and Gynaecologists (ACOG). Continued buprenorphine treatment is safe for women who become pregnant while already in this type of treatment. It achieves similar outcomes to methadone in reducing maternal drug use and promotes positive neonatal outcomes, and appears to result in a less severe neonatal abstinence syndrome (Young & Martin, Jones et al. 2012). However as buprenorphine induction may place the foetus at risk from precipitated maternal withdrawal and potentially trigger relapse in mothers, methadone is recommended for active opioid users who need to be inducted to pharmacotherapy during pregnancy (Young & Martin 2012).



Women's AOD Services Network members, who were consulted in the development of this resource, reported incidents involving case workers from Family and Community Services (FaCS) who had stipulated a reduction in, or cessation of, pharmacotherapy as a condition for either retaining a child in the mother's care or for restoration of the child to her care. Major concerns were raised over these types of conditions as they are inconsistent with evidence-based medical practice of pregnant women with an opioid dependence. Cessation of pharmacotherapy is a common trigger for relapse to illicit drug use, which is inconsistent with the goal of a safe and stable family environment for children and is an undesirable outcome for the mother. Improving communication and building understanding within FaCS may assist to improve outcomes in this regard.

FaCS will [sometimes] dictate treatment; they'll say "mum needs to come off methadone" which makes it really unsafe for her and her children... the methadone or Suboxone is actually a stabilising factor in their lives...

Sometimes the client doesn't want to come off and they know how unstable they'll become, but they sometimes get it into their mind... and will actively try and get the prescriber to bring them down in the community which is even more unsafe... all driven to get their child back in their care.

We have had a few women who have been on prescribed Valium [in conjunction with pharmacotherapy] for however long and it's like "bang, you need to be off it"... and we've had a couple of hairy [challenging] situations where that's been the expectation and we've had to try to explain again [to the FACS worker] the slow [withdrawal] process that needs to occur.

# 1.4 Women accessing treatment for a significant other's drug use

When it comes to accessing AOD treatment for someone else's substance use, women in Australia are twice as likely as men to have sought assistance (AIHW 2014). Women may access AOD treatment to assist them to find answers as to why a loved one is using alcohol and other drugs and how they can receive support. An empathetic and non-judgemental approach is essential in order to engage women who may feel isolated from the AOD treatment sector because their partners are the focus of care and treatment.

"Blame, guilt, grief, shame and frustration are natural companions of the trauma of AOD problems and other major family difficulties in our culture–families have needs in their own right and have a right to have their needs acknowledged" (Marsh et al. 2013)."

Support and a sense of being heard can help significant others to manage their concerns and reduce levels of anxiety. Copello and colleagues (2005) recommend the following five steps when working with a family member or partner who is living with someone with AOD-related problems. These steps aim to reduce the level of strain (physical and psychological) experienced by the family member or partner, and enhance their coping mechanisms:

- Give the family member or partner the opportunity to talk about the problem and be heard.
- Provide relevant information in accordance with the needs of the client.
- Explore how the family member or partner responds to the person's substance use.
- Explore ways of enhancing social support.
- Discuss the possibility of onward referral for further specialist help.

"Practice that is family inclusive does not require specialist family therapy training and can result in family members receiving the support they need in their own right and can also be beneficial to treatment outcomes for the drug user. However, if counsellors wish to engage in family therapy, specific skills and specialist training is required" (Marsh et al. 2013).



**Find out more...**about supports for women who attend AOD treatment for another person's problematic substance use.

#### Family Drug Support

Family Drug Support (FDS) was formed in 1997 after its founder, Tony Trimingham lost his son to a heroin overdose. FDS is a caring, non-religious and non-judgemental organisation, primarily made up of volunteers who have experienced firsthand the trauma and chaos of having family members with drug dependency. FDS provides a 24 hour/7 day a week telephone support line, website, support groups and courses for family members and significant others. Website: www.fds.org.au

#### Tools for Change: A new way of working with families and carers. NADA 2009

This resource seeks to improve the support offered to the families and carers of clients who access AOD services in NSW, moving towards a more family inclusive model where all parties are considered engaged, supported and involved.

#### A Checklist for Family Sensitive Practice for the Alcohol and Other Drug Field. NCETA 2010

This checklist comes as part of the Family Sensitive Policy and Practice Toolkit, and provides a comprehensive checklist for reviewing how oriented to family inclusive practice an AOD service is.

Website: www.nceta.flinders.edu.au

#### Single Session Therapy

Single Session Work has a range of applications and can be utilised by services which primarily work with individuals to engage family members in the 'treatment' process. This targeted approach is appealing to families because it is often difficult for family members to commit to ongoing work. A surprising amount can be achieved in one or two sessions and the SSW approach always leaves the 'door open' for further work with the family if required.

Website: www.bouverie.org.au

# CREATING AN ENABLING ENVIRONMENT FOR WOMEN: BEST PRACTICE APPROACHES FOR AOD TREATMENT



- An enabling environment for women requires a gender-responsive approach where AOD treatment is shaped by, and is responsive to, women and their experiences.
- A strengths-based approach to AOD treatment with women focuses on their strengths and resilience, contributing to capacity building and client self-determination.
- **Trauma informed service provision** acknowledges the lived experience of trauma common to women and is sensitive to avoiding perpetuating traumatic events.
- Family inclusive practice responds to the significance of familial relationships for women. Strengths can be harnessed among families or within communities to support positive and sustained treatment outcomes.
- Evidence-based interventions for AOD treatment such as behaviour therapies, cognitive behaviour therapies, motivational interviewing and 12-step groups are suitable for women. Good practice insists on comprehensive assessment and individualised treatment planning.
- Coordinated case management and integrated programs benefit women and overcomes fragmented service systems.
- **Sexual safety and sexual health** are part of a comprehensive AOD treatment plan for women accessing AOD treatment.
- Assertive follow up, outreach and aftercare are interventions applied in the AOD treatment sector that assist in engaging and retaining women in treatment long enough to establish effective and sustained improvements to their lives.

All the treatment approaches explored in this resource have consistent and complementary messages. They are easily adaptable to the AOD treatment sector, and have been implemented in Australia and overseas contributing to a solid evidence base.

The following six components of women's treatment have been identified as promoting positive treatment outcomes: access to childcare; access to prenatal care; access to women-only programs; psychoeducational sessions focused on women-specific topics; mental health interventions and comprehensive services that offer multiple components.

Positive treatment outcomes from the components described above are longer retention in treatment; decreased substance use; decreased mental health symptoms; improved birth outcomes; greater employment; HIV risk reduction and improved self-reported health status (Ashley et al. 2003).

The best practice interventions identified in this resource have relevance to, and are able to be adapted in, mixed-gender services and specialist women's AOD services. In this section various approaches and interventions are presented, with reference to comprehensive practice guideline documents for a more expansive description and implementation strategies.

#### IMPROVING ACCESS

Improving access for women into AOD treatment by being responsive to the needs and issues they present with requires some flexibility and planning – facilitated by engagement skills and comprehensive assessment. Even if the result is to provide support and referral onto a specialist service, it is essential that workers and treatment services enlist the skills they have to engage women and encourage them to take the first step into treatment.



-rom the frontline

Most women indicated that gaining access to treatment without waiting made it easier to receive help. Other treatment enablers were:

- low or free cost of treatment
- a suitable location
- access to childcare
- access to transport
- a women-only service

When I went to [service] they got me in straight away. There were [also] local aftercare groups- very helpful.

I only waited two weeks. It was really easy to get into and the price is really good, I think a lot cheaper than the last rehab I was at.

#### EFFECTIVE INTERVENTIONS WHEN WORKING WITH WOMEN

The literature tells us that effective women specific interventions include:

- family-inclusive practices that focus on repairing relationships with children and family members and enhancing the quality of the family/domestic environment
- addressing trauma
- developing support systems to prevent relapse
- comprehensive service models for pregnant women
- parenting skills for mothers on methadone [and other pharmacotherapy] maintenance
- relapse prevention for women with PTSD, marital distress and alcohol dependence
- dialectical behaviour therapy for patients with co-occurring drug dependence and borderline personality disorder. (Greenfield et al. 2007).

In a longitudinal study of recovery and relapse cycles, self-help participation was more strongly associated with moving from substance use to recovery for women than for men in substance use treatment, and women were more likely than men to participate in self-help (Grella et al. 2008).

This section provides an overview of a number of the approaches and interventions outlined above with reference to source documents that provide more comprehensive articulation of each topic. The case study over the page also provides an example of how a number of these interventions can work together.



**Find out more...**about engagement and assessment processes.

# NSW Health Drug and Alcohol Psychosocial Interventions: Professional Practice Guidelines NSW Health 2008

These guidelines aim to provide a benchmark for the delivery of quality psychosocial interventions within AOD treatment services. They recognise the value of such interventions within the AOD field, and support professional implementation of them. They emphasise the need for better understanding about the purpose and benefits of the interventions.



#### Case study

Kylie is a 22 year old Aboriginal woman referred to our service by Aboriginal Legal Aid (ALS) for detox and rehab for poly drug dependence. She first used illicit substances at age 13 and disclosed daily methamphetamine and cannabis use, and regular heroin use. Prior to referral, Kylie had presented to the local antenatal service at 27 weeks pregnant and had not previously attended antenatal appointments. She was engaged by the Chemical Use in Pregnancy Service (CUPS) team at the local hospital and allocated a Family and Community Services (FaCS) caseworker. Kylie was involved with the criminal justice system and had current charges for which the diversion of incarceration was a motivating factor to seek treatment. She reported a previous suicide attempt and a history of depression.

Kylie is exposed to domestic violence in her current relationship, and while in treatment sought an Apprehended Violence Order (AVO) to protect herself and her unborn child. Kylie's parents have a history of alcohol dependence and while she had resided in her parents' home prior to admission, FaCs deemed it unsuitable for her to return there with her newborn child. Despite these stressors, Kylie was willing to engage in treatment and demonstrated insight into the need to provide a safe environment for herself and her unborn child. She worked openly with staff and external formal supports such as FaCS, CUPS and the ALS to improve her circumstances.

The goal of treatment was for Kylie to detox from illicit substances prior to the birth of her child. Treatment also provided the opportunity for her to stabilise socially and develop a safe and supportive long-term plan. By remaining with us and participating in our short term rehabilitation program following the birth, staff were able to assist Kylie in her transition to motherhood and provide coaching on parenting skills. FaCS recommended that she attend a longer-term rehabilitation program; however, Kylie was able to negotiate an outpatient support program so that she could settle back into the community with her baby. On completion of the program she returned to the community, where she was supported by numerous services to secure appropriate accommodation and engage in key AOD aftercare and parenting supports. Being flexible and extending the service to accommodate the unique needs of Kylie in this circumstance was of great benefit to the client and her family.

Kylie's case study outlines not only the concurrent issues that required attention alongside her substance use, but the benefits of applying a number of best practice approaches for women, including: coordinated case management; harnessing of specialist services not able to be provided by the AOD treatment service; attending to issues related to FDV and mental health issues; being trauma informed; and allowing for client self-determination.

**Counselling guidelines: Alcohol and other drug issues** 3rd Edition WA Drug and Alcohol Authority (Marsh et al. 2013)

These guidelines includes over 50 chapters exploring some of the key skills needed to work at an individual level with people who have AOD problems including a case management, group work, referral, working with intoxicated and incarcerated clients, clinical supervision and stress and burnout.

Talking therapies for people with problematic substance use Te Pou o Te Whakaaro Nui 2010

This guide provides useful background information on problematic substance use, suggestions for interventions during the various stages of treatment, and information on the range of talking therapies currently utilised by practitioners in New Zealand. Also includes sections on Maori, other Pacific Islander cultures and working with refugees.

#### 2.1 Establishing a gender-responsive service

A gender-responsive approach attends to the experiences of women and, in the context of AOD treatment, focuses on how substance use and dependence is shaped by gender. It also explores how the socialisation process can impact treatment experiences. Being gender-responsive means that the planning of AOD interventions, the language used, how it is delivered and by who, is explored and developed according to the experiences of women. In this context, gender is understood to be "what a society believes about the appropriate roles, duties, rights, responsibilities, accepted behaviours, opportunities and status of women and men in relation to one another. In simple words, people are born female or male but learn to be girls and boys who grow into women and men. This learnt behaviour makes up gender identity and determines gender roles." (WHO 2009).

"Being gender-responsive means that the planning of AOD interventions, the language used, how it is delivered and by who, is explored and developed according to the experiences of women"

Gender-responsive services are those that aim to understand and address issues experienced by women. Gender-responsive approaches may include women-only programs and groups; services with only female staff; provision of on-site childcare; pre-natal, parenting, or other child-related and gender-sensitive program content; and, therapies that focus on issues prevalent among women such as trauma from physical, emotional, and sexual abuse (Covington 2008; Greenfield 2009, Prendergast et al. 2011). While all of these interventions may not be possible in a mixed-gender service at all times, there are areas that might be adapted – such as women-only groups and mixed-gender groups that have very specific guidelines around sensitivity to gender issues.

There is an opportunity to apply a gender-responsive approach in a mixed-gender service that leads to positive outcomes for men such as improved awareness of negative attitudes towards women and their effects. More specifically, through male staff involvement in some aspects of gender-responsive approaches, there can be transfer of learning and role modelling. This approach can have the dual effect of educating male clients in improved behaviours regarding women and reinforcing supportive, non-abusive relationships between men and women. It is important to remember that describing an AOD treatment service as being gender-responsive means that there is more within programs and interventions than simply raising awareness it means that there is a commitment to actually doing something about gender inequality.

Research findings are inconsistent as to whether women-only AOD treatment is more effective for women compared to mixed-gender services. However, incorporating gender-responsive practices within a mixed-gender service could enhance access and retention for women. **Section 6.1** explores specific ways of reviewing practices at a mixed-gender service in order to make it more gender-responsive.



**Find out more...**about gender-responsive programming.

# Integrating Gender into HIV/AIDS Programmes in the Health Sector: Tool to Improve Responsiveness to Women's Needs. WHO 2009

This tool helps health-care providers integrate gender into HIV/AIDS programmes they wish to set up, implement and evaluate so they are more responsive to women's needs. It suggests practical actions to address key gender issues. Although related to HIV services the key messages regarding gender responsiveness are easily translated into the AOD sector.

Gender-Responsive Treatment for Women with Substance Use Disorders WATCH (Women and Their Children Heal) 2007 (Bastek-Karasaw & Davis)

The purpose of this document is to offer agencies and programs that serve women with substance use disorders across the continuum of care, a set of sample standards and recommendations for treating problematic drug use in women throughout their lifespan. These recommendations serve as a guide for creating quality gender-responsive treatment and support services for women within a biopsychosocial context, ultimately enhancing their outcomes. The information and recommendations presented here are based on research, reflect best practices in the treatment of addiction in women, and are grounded in women's experiences.

# 2.2 Adopting a strengths-based approach to treatment provision

A strengths-based approach focuses on finding strengths within the individual, their networks and community. Examples of strengths might be personal competencies, resources, motivations, personal characteristics and positive relationships (McCashen, 2005; Saleeby, 1996). Working from a strengths perspective means identifying positives within individuals and building their network/environment strengths as opposed to the traditional deficits and/or pathology perspective. The strengths-based approach is compatible with resilience theory where the central principle is that in spite of facing adversity people often do well and thrive (Masten 2001) and also encompasses other theories and broader ideas such as empowerment and wellness (Saleeby, 1996).

A strengths-based approach seeks to reveal and reinforce the strengths women bring with them to AOD treatment. Simple ways of including this approach to working with women in AOD services can be through strengths-based questioning. Here are some examples:

- What did it take for you to be here today?
- What is working well for you?
- What have you tried, and what has been helpful in making changes?
- You are resilient, what do you think helps you bounce back?
- What would things look like if they were working better for you?

(Adapted from Change Lives, New Journeys 2013).

A strengths-based approach places women at the centre of AOD treatment provision as the experts on their lives. With this core value underpinning AOD interventions, treatment occurs in a manner that values the experiences of women and seeks to gather the support of willing collaborators that could include other support services, families and community groups. As with a gender-responsive approach, a strengths-based approach to AOD treatment underpins and informs all proposed interventions in ways such as thinking about how language is used, how programs are run and ensuring client consultation. It is embedded in the way organisation themselves are managed.



Find out more...about a strengths-based approach.

#### Strengths-Based Practice: The Evidence UnitingCare (Scerra 2011)

This discussion paper explores the evidence of strengths-based practice, and reports how strengths-based principles can be enacted in different ways in a range of contexts to achieve positive outcomes and to empower the children, young people and families.

#### Change Lives, New Journeys

A website that relates to living with and supporting those with a lived experience of brain injury – adopting a strengths-based approach.

Website: www.changedlivesnewjourneys.com

#### 2.3 Trauma informed care and trauma specific approaches

Trauma is often characterised as resulting from a one-off event. Yet repeated extreme interpersonal trauma resulting from adverse childhood events ('complex' trauma) is not only more common, but far more prevalent than currently acknowledged, even within the mental health sector. The effects of complex (cumulative, underlying) trauma are pervasive and, if left unresolved, can negatively impact the mental and physical health across a person's lifespan. Research shows that the impacts of even severe early trauma can be resolved, and its negative intergenerational effects can be intercepted. People can and do recover and their children can do well. For this to occur, mental health and human service delivery needs to reflect current research insights (Kezelman & Stavropoulos 2012).

AOD treatment services and workers wanting to become trauma informed and engage in trauma-informed care need first connect with these five core principles (adapted from Complex Needs Capable, NADA 2013, based on ASCA practice guidelines for trauma informed care):

- **Safety** How can workers/AOD services be changed to ensure physical and emotional safety more effectively and consistently?
- **Trustworthiness** How might a service modify practices so as to engender trustworthiness through task clarity, consistency, transparency and interpersonal boundaries?
- **Choice** How can services be modified to ensure consumer experiences of choice and control are maximised?
- **Collaboration** How can services be modified to ensure that collaboration and power-sharing are maximised?
- **Empowerment** How can services be modified to ensure that experiences of empowerment and the development or enhancement of client skills are maximised?

Women with substance use disorders, and especially those with co-occurring mental health disorders, have a higher likelihood of past experience of traumatic events, including sexual assault and domestic violence.

Drug and alcohol use disorders, mental health and trauma are now recognised as critical and interrelated issues, requiring comprehensive treatment responses (NIDA 2009). Factors associated with early attrition among women with co-occurring PTSD and substance use disorders in treatment include experience of partner violence during adolescence (McCrady et al. 2009). The addition of trauma-focused group behavioural treatments to existing substance abuse treatment has been found to reduce PTSD symptoms in women with co-occurring substance disorders (Resko & Mendoza 2012).

Study findings indicate that despite concerns about potential negative impact of addressing trauma in an AOD context, positive outcomes have been achieved and the interventions have been successfully delivered by a range of clinicians, not necessarily just those with higher degrees or advanced expertise in the area of trauma (Bates et al. 2002).

Trauma informed practice is highly congruent with evidence-based AOD intervention, as outlined in the following table adapted from Trauma Matters: Informed Practices in Women's Substance Use Services (The Jean Tweed Centre 2013).

# Examples of Congruence Between Trauma-Informed Practices and Evidence-Based Substance Use Approaches

TRAUMA INFORMED PRACTICE	EVIDENCE-BASED SUBSTANCE USE APPROACHES
Pacing – avoiding the rush to disclosure (the 'action stage' of the trauma recovery experience) and accepting that a woman need not necessarily disclose her experience of trauma	Stages of Change – recognising that different strategies are necessary based on the woman's stage of change regarding her substance use and that action strategies are only appropriate in the action stage of change
Relationship building – accepting the challenge to construct a therapeutic connection over time, recognising that trustworthiness is built slowly and respectfully by honouring the woman's background and experiences	Harm reduction principles – finding goals that are consistent with the woman's current abilities and situation, not necessarily requiring abstinence as a goal
Ensuring safety in the counselling relationship – avoiding aggressively confrontational approaches that can re-traumatise and obstruct safety	Motivational Interviewing – avoiding shame, blame and guilt and focusing instead on learning from setbacks
Providing choice and control	<b>Motivational Interviewing</b> – internalising the focus of control and highlighting self-efficacy
Building empowerment skills	Motivational Interviewing – building on the woman's strengths; asking questions that elicit information, rather than telling in an instructional way
Using a collaborative approach with women instead of a top-down hierarchical model	Harm reduction principles – taking small steps approach
	Relationship building – the quality of interpersonal relationship may determine whether or not women remain engaged in the process of change; the quality of the interpersonal relationships may be more important that the concrete services received
Strengths and skills based	Motivational Interviewing – providing a menu of options that respond to concurrent life circumstances; recognising the need for support that address a range of areas, such as food security, child care, adequate housing and poverty

The key message is that trauma informed care can be implemented in any service setting and is consistent with strengths-based approaches and family inclusive practices, both of which place the client and worker in a collaborative partnership.

Comprehensive services integrating mental health, substance use and trauma-informed services may provide more effective treatment for women than non-integrated service settings. Provision of more integrated counselling (addressing trauma, mental health and substance use issues) is associated with better outcomes for women (Greenfield 2009).

While the nature of trauma is complex and challenging, the variety of supportive resources and toolkits is vast. A commitment to learn, train and educate around trauma-informed practice is a demonstration of support from AOD workers to women who experience trauma, serving to enhance engagement and retention in AOD treatment.





The impact of enduring symptoms of trauma on AOD treatment and women's quality of life were highlighted during consultations held with AOD treatment services who informed this resource. Some of the issues thought to be a consequence of past trauma that were raised included self-harming behaviours, emotion dysregulation, difficulties with interpersonal relationships, recognising and/or maintaining healthy boundaries, and vulnerability to future trauma. As examples, several informants highlighted the difficulties faced by women who begin emotionally charged romantic relationships during residential treatment, and the negative effect the relationships had on their treatment outcomes when the relationships failed.

"the message from the frontline is that a good understanding of trauma and its ongoing effects on women and their children is essential for all workers in AOD treatment settings."

The increased risk of trauma among Aboriginal women, including intergenerational trauma, was also noted by some informants. AOD workers also raised the issue of the trauma experienced by children following separation from their mothers as a result of Care and Protection Orders, and the long-term implications of such trauma on children's mental health. Network members suggested that opportunities be explored for providing mental health care for children in the context of drug and alcohol treatment for mothers. Overall, the message from the frontline is that a good understanding of trauma and its ongoing effects on women and their children is essential for all workers in AOD treatment settings.

Child sexual abuse, adult sexual assault, high levels of domestic violence and children being removed... we have significant issues with women relating to others and lacking emotion regulation.

In my experience [the issue of trauma] has been substantial. I can't really think of a woman in drug treatment who doesn't have a complex trauma history.

I'd say 90% of the young women we see have a history of some type of abuse, or they've witnessed traumatic events. Many of them come to us with self-harming behaviours, personality disorders, PTSD... It can be difficult because we're a mixed gender service and they're vulnerable to forming a volatile relationship while they're with us.

We help women focus on what is a healthy relationship... If domestic violence isn't addressed women will continue to make unhealthy relationship choices.



Find out more...about trauma informed treatment.

# Trauma Matters: Guidelines for Trauma-Informed Practices in Women's Substance Use Services Canadian Women's Health Network 2013

The guidelines set out in this document focus specifically on the intersections of trauma and substance use issues among adult women.

# Trauma-Informed Treatment Guide for Working with Women with Alcohol and Other Drug Issues 2nd Edition (Marsh et al. 2012)

This manual is for service providers working with women experiencing alcohol and other drug and mental health issues. The manual covers a wide range of topics, such as: assessment and screening; case formulation and the neurobiology of trauma.

Information is also included on working with different populations, such as Aboriginal women and culturally and linguistically diverse women. The manual takes a practical approach with a 'tip sheet' at the end of each chapter and a number of useful appendices, including: information sheets for clients on a range of topics; information for workers to consider when confronting a number of issues, such as domestic violence and assessment tools.

#### Trauma-Informed Organizational Toolkit for homelessness services (Guarino et al. 2009)

This toolkit offers homeless service providers with concrete guidelines for how to modify their practices and policies to ensure that they are responding appropriately to the needs of families who have experienced traumatic stress. AOD services can also use this resource to assist them in becoming trauma informed.

#### 2.4 Family-inclusive practice

A family inclusive or family sensitive approach, recognises the possibility of harnessing constructive support from family members. It's worth noting that the approach acknowledges that the meaning of 'family' is defined by the client and can include significant others and carers. As already outlined in the previous section of this resource, family and family life for women can be particularly significant and it is therefore a useful target for intervention.

AOD workers have an important role to play in harnessing family inclusivity as a support for treatment with women. According to the Bouverie Centre, "All workers are capable of generating practical ideas for the way your service operates that will improve the quality of life for family, carers, clients and other workers." (1998).

A family inclusive practice approach recognises that interventions are more effective when they include family members. It supports the view that individuals influence other members in their environment, especially family, and that family members, in turn, have an impact on these individuals (Kina Family and Addictions Trust 2005, in NADA 2009). The Tools for Change toolkit was specifically designed to inform AOD workers about working better with family members and carers of problematic substance users was produced by NADA (2009), and provides practical tips, interventions, resources and examples of AOD organisations that have implemented a family inclusive practice approach. This is a good place to start when considering how family inclusive practice might be applied to the work you do with clients.

The NADA toolkit provides interventions that address three broad categories as identified by Copello, Vellman and Templeton:

- Working with family members to promote the engagement of clients into services
- Involving families and others in treatment plans
- Providing services directly for family members (2005).

The family inclusive practice approach also attends specifically to parenting. While parenting is not the exclusive preserve of women, support for improved parenting has benefits for both mother and child(ren). "Women are typically primary carers, especially in single parent families," (Marsh et al. 2012) and while being a parent can be a potential barrier to women accessing AOD treatment, it can be a motivating factor for reaching out and getting support. Family inclusive practices seek to build on the strengths of the parent-child relationship while working with areas of difficulty. Specific parenting related interventions are explored in **Section 4**.



I heard [service] has a good reputation, it is a long-term rehab which I need. I liked the sound of the program and that my son could visit and stay. It's also very family orientated and [caters to] a smaller number of women.

I picked this service because it deals with the issues that I need to deal with, such as parenting issues.

While many services may not have the option for children to stay or be able to run dedicated parenting programs, there are alternative strategies they can utilise for being more family inclusive, as outlined below:

- Setting up specific times and appropriate spaces for family/children to visit.
- Having particular environments that are welcoming to children; eg, waiting areas or gardens.
- Involving specialist parenting services to provide workshops to clients as part of their regular program.
- Using parenting examples in the context of therapeutic groups as appropriate many of the skills are relevant to 'coping skills rehearsals' and identifying the stressors that might be a precursor to relapse.
- Engaging clients in developing information brochures that target family members, providing details about the experience of AOD treatment as a way of demystifying the process.
- Having open days, art exhibitions, picnics and other activities so that family members can attend.
- Single session therapy, an effective and efficient therapeutic approach that addresses family needs without requiring a long-term commitment to therapy.

Family inclusive approaches start with conversations about families and significant others. Within the context of an assessment, questions related to family, children and parenting can be asked that provide some acknowledgment of the impact families and significant others may be having and furthermore, what plans might be made to put in place to be inclusive of them. For more concrete examples of incorporating family inclusive practice in your workplace refer to **Section 6** of this resource.



**Find out more...** about Family Inclusive and Family Sensitive Practice.

#### Tools for Change: A new way of working with families and carers NADA 2009

This toolkit seeks to improve the support offered to the families and carers of clients who access AOD services in NSW, moving towards a more family inclusive model where all parties are considered engaged, supported and involved.

#### A Checklist for Family Sensitive Practice for the Alcohol and Other Drug Field NCETA 2010

This checklist comes as part of the Family Sensitive Policy and Practice Toolkit, and provides a comprehensive checklist for reviewing how oriented to family inclusive practice an AOD service is.

# 2.5 Evidence based interventions and coordinated case management

The approaches outlined above (gender-responsive, strengths-based, trauma informed and family inclusive) provide best practice frameworks to AOD treatment for women. These frameworks guide and shape the way a range of therapeutic modalities, accepted as effective for use with general AOD treatment populations, can be applied when working with women. In short, many of these generalist interventions are aligned with the needs of female substance users or are capable of being adapted to gender-specific treatment needs, including:

• **Behavioural therapies:** Dialectical Behaviour Therapy (DBT), Acceptance and Commitment Therapy (ACT), Community Reinforcement Approach (CRA)

These therapies focus on improving interpersonal, self-regulation and distress tolerance skills by integrating behaviour strategies and mindfulness practices. CRA for example seeks to reinforce positive or non-substance using behaviours as opposed to ongoing substance use behaviours.

#### Cognitive Behavioural Therapy (CBT)

CBT is a structured therapy that aims at adjusting thoughts and behaviours that control problematic behaviours. CBT focuses on the cognitions, or interpretations of events that lead to actions and emotions incorporating strategies such as skills rehearsals and relapse prevention.

#### Motivational Interviewing (MI)

Motivational Interviewing is based on the Stages of Change (Prochaska & DiClemente 1982), and incorporates a specific set of interviewing strategies designed to guide an individual through different stages in the process of changing behaviour. The core goals of MI are to generate behaviour change by assisting the individual to resolve ambivalence about treatment and reduce their substance use.

#### • Narrative Therapy and Mindfulness

Narrative Therapy is a collaborative and non-pathologising approach to counselling and community work which centres people as the experts of their own lives. A narrative approach views problems as separate from people and assumes people as having many skills, abilities, values, commitments, beliefs and competencies that will assist them to change their relationship with the problems influencing their lives. It is a way of working that considers the broader context of people's lives particularly in the various dimensions of diversity including class, race, gender, sexual orientation and ability.

Mindfulness is a modern reworking of ancient meditation traditions, principally Buddhist. It is designed to help people deal with day to day difficulties by putting the individual in control of their own mind. The aim of mindfulness therapy is to help the individual learn to be aware of their thoughts and bodily sensations and in so doing be able to better cope with day to day emotions and problems.

While these approaches are less researched as those described above, they may be just as effective and are in keeping with the themes of trauma-informed practice, a strength-based approach, a gender-responsive approach and family inclusive practice.

#### Self-help groups

Self-help groups are groups of people with common experiences related to their substance use that meet together in order to support each other, share practical help and obtain goals of either abstinence or reduced harm associated with ongoing substance use.

#### • Peer support work

Peer support work is undertaken by those who have a lived experience of substance use and/ or have other common experiences such as their age or social connections. Peer support involves a range of activities designed to support, in a non-professional way, others who may be distressed or isolated. When adapting these treatment modalities to match with the frameworks already explored as best practice in the AOD treatment of women, the following questions can be asked:

- 1. How might these therapeutic modalities be shaped by gender?
- 2. What modifications in language, approach or delivery might make them more gender sensitive?
- 3. How might the intervention be tailored for specific client needs?

For interventions to be most effective, they need to be tailored to clients' needs and address specific risk factors. The evidence indicates that the overarching principles of what works in women's treatment for drug and alcohol issues are targeted and gender responsive services. Furthermore, AOD treatment service workers are often well placed to coordinate or be part of a coordinated case management approach that ensures services 'wrap around' the women they are trying to support. Exploration of coordinated case management, as adapted from the Complex Needs Capable resource (NADA 2013), is presented in the next section.



**Find out more...** about different therapeutic modalities outlined above.

# NSW Health Drug and Alcohol Psychosocial Interventions: Professional Practice Guidelines NSW Health 2008

These guidelines aim to provide a benchmark for the delivery of quality psychosocial interventions within AOD treatment services. They recognise the value of such interventions within the AOD field, and support professional implementation of them. They emphasise the need for better understanding about the purpose and benefits of the interventions.

**Counselling guidelines: Alcohol and other drug issues** 3rd Edition WA Drug and Alcohol Authority (Marsh et al. 2013)

These guidelines includes over 50 chapters exploring some of the key skills needed to work at an individual level with people who have AOD problems including a case management, group work, referral, working with intoxicated and incarcerated clients, clinical supervision and stress and burnout.

**Talking Therapies for people with Problematic Substance Use** Te Pou o Te Whakaaro Nui 2010

This guide provides very useful background information on problematic substance use, suggestions for interventions during the various stages of treatment, and information on the range of talking therapies currently utilised by practitioners in New Zealand. Also includes sections on Maori, other Pacific Islander cultures and working with refugees.

# COORDINATED CASE MANAGEMENT

Case management is a process designed to coordinate supports, interventions and other services in response to the needs identified by clients through assessment. One of the important features of the coordinated case management of women engaged in AOD treatment is a collaborative approach that places the client at the centre of the planning. In this approach, the AOD worker acts very much like a coordinator of both the therapeutic interventions they can provide given their skill base, as well as the sourcing and implementation of external specialist services. It is useful to seek clarity within your organisation about identifying a particular approach to coordinated case management given that different approaches have different levels of intensity (NSW Dept. of Health 2008).

As outlined in the NADA Complex Needs Capable resource, there are two preferred case management approaches when working alongside clients with complex needs: Proactive and Assertive case management (2013), and these are in keeping with the needs of women in AOD treatment. The following elements are essential whichever case management approach is implemented:

- Establishing a trusting relationship with clients
- Establish close and frequent communication with them (eg, check-ins and regularly scheduled meetings)
- Develop a care plan based on their goals, strengths and needs
- Facilitate transportation for important appointments
- Liaise with other service providers with consent
- Assist the development of life skills and/or the person's participation in healthy, safe activities
- Establish close communication with supportive significant others in the person's life
- Advocate for improved communication among service providers, continuity of care and access to care/services.

(Complex Needs Capable, NADA 2013:35)



**Find out more...** about various types of case management.

#### Complex Needs Capable: A Practice Resource for Drug and Alcohol Services NADA 2013

Complex Needs Capable and the accompanying website was developed to build capacity within the non-government drug and alcohol sector by supporting the development of staff skills, knowledge and confidence, and of organisational capacity in responding to the needs of people with drug and alcohol issues and complex needs such as cognitive impairment and criminal justice system contact.

Website: www.complexneedscapable.org.au

#### Case management in the substance abuse field: insights from the research literature 2005 (McDonald)

Prepared as a contribution to the process of establishing new policies and activities in case management, this paper presents insights from the published research literature, identifying and assessing how case management is used in the AOD sector and its results.

#### NSW Health Drug and Alcohol Psychosocial Interventions: Professional Practice Guidelines NSW Health 2008

These guidelines aim to provide a benchmark for the delivery of quality psychosocial interventions within AOD treatment services. They recognise the value of such interventions within the AOD field, and support professional implementation of them. They emphasise the need for better understanding about the purpose and benefits of the interventions.

**Counselling guidelines: Alcohol and other drug issues** 3rd Edition WA Drug and Alcohol Authority (Marsh et al. 2013)

These guidelines includes over 50 chapters exploring some of the key skills needed to work at an individual level with people who have AOD problems including a case management, group work, referral, working with intoxicated and incarcerated clients, clinical supervision and stress and burnout.

#### Pathways of recovery 2005 (Rickwood)

The paper considers definitions of the terms 'relapse' and 'relapse prevention' and discusses the place of relapse prevention within the spectrum of interventions for mental health. Relapse prevention and the related concept of rehabilitation are placed under the umbrella of recovery, which comprises treatment and continuing care.

#### 2.6 Women's sexual health and safety in AOD treatment

Substance use is frequently a gendered experience for women, where initial substance use experimentation can be associated with intimate relationships and sexual experiences. While the exploration of links between substance use and sexual experiences is the domain of therapeutic work in AOD treatment, there are opportunities to provide information and education specifically around sexual and reproductive health, and contraception in particular.

"Negotiating safe sex and contraceptive use can be difficult for young women (WHO, 2006) even when they can access contraception. Power differentials in relationships mean they are not always in a position to insist on condom use, and young women with poor body image or those in violent relationships are more at risk. These types of findings reinforce the importance of exploring sexual behaviour and safety whenever women present for assistance in an AOD treatment setting. Furthermore, there are strong associations between women's problematic substance use and traumatic sexual events, such as childhood sexual abuse and sexual assault as an adult (Willis & Rushforth 2003). Substance use is a common coping mechanism for dealing with sexual trauma.

Women who have lengthy histories of problematic substance use may be less likely to have engaged with routine sexual and reproductive health check-ups; these areas of need may be addressed in the context of AOD treatment via referral to a specialist clinic. Furthermore, AOD treatment provides an opportunity to engage in harm minimisation education about STIs as well as BBVs (hepatitis C and HIV).

The gendered experience of drug use also extends to injecting behaviour where women are frequently introduced to injecting drug use by male partners who may then control the attainment of drugs and injecting equipment (Pinkham & Sempruch 2007; Shannon et al. 2008). Women have been found to be more likely than men to use drugs with many partners; share paraphernalia after a partner has injected; have difficulty negotiating condom use with their sex partners; and, exchange sex for money or drugs (Maher & Hudson 2007; Ashley et al. 2009)

The prevalence of women entering treatment who have in the past year exchanged sex for money and/or drugs is high (30-40%) (Burnette et al. 2008; Sallmann 2010). Women who work in the sex industry, particularly street-based sex workers, have high rates of drug and alcohol use and dependence, including intravenous drug use and poly-drug use, as well as increased risk of contracting blood borne viruses (Roxburgh 2008; Simpson & McNulty 2008). A recent study found that women who work in the illegal sex industry have significantly poorer mental health than those working in brothels or privately (Seib et al. 2009).

An investigation into street-based sex work in Sydney found that many workers initiated sex work to obtain money for drugs (Roxburgh 2008). Working in the sex industry is associated with higher risk of physical and mental health problems, primarily among those involved in street-based sex work, and is often linked with socio-demographic disadvantage.

Harm reduction and outreach are indicated as best practice approaches for women who engage in street-based sex work, given their transience. However, AOD treatment services that use best practice modalities (as outlined in **Section 2.6**) that are gender-responsive and trauma informed are also effective.

#### ESTABLISHING AN ENVIRONMENT THAT PROMOTES SEXUAL SAFETY

Having awareness of the impact of past experiences of sexual assault on women that present for support to AOD treatment services, is essential to ensure that future sexual safety be a consideration – particularly in mixed-gender residential services. The following aspects of AOD service provision should be considered in order to promote sexual safety.

- Review the environment by specifically taking into account sexual safety.
- Review policies and procedures that relate to how staff and clients use the physical spaces.
- Integrate trauma informed care organisation-wide and to all aspects of treatment.
- Clearly display education material regarding sexual safety.
- Provide education to all clients about sexual safety and their responsibilities regarding the sexual safety of others.
- Ensure there are policies and procedures regarding response to alleged sexual assault.
- Make links with specialist services regarding sexual safety and sexual trauma.
- Availability of condoms, dental dams and lube.

For information on safety for women in a mixed-gender setting see Section 6.1.



Find out more...about sexual safety (examples taken from the mental health sector).

#### Sexual Safety of Mental Health Consumers Guidelines NSW Health 2013

The purpose of these guidelines is to support mental health services to meet their responsibilities in relation to the sexual safety of mental health consumers through the provision of practical advice and strategies; promote sexual safety to key stakeholders, health staff and managers, consumers and their families and carers; clearly outline the information consumers and their families and carers should receive about their rights and obligations in relation to the sexual safety of consumers; and improve collaboration and strengthen relationships between mental health services and sexual assault services.

#### **Promoting Sexual Safety, responding to sexual activity, and managing allegations** Vic. Health 2009

This Chief Psychiatrist's guideline sets out the relevant legislation and policy, and establishes minimum standards for the clinical management of sexual activity, and the appropriate staff response to patients who report inappropriate sexual activity in adult acute inpatient units. It provides guidance on decision making processes and clarifies the circumstances where clinical discretion may apply.



**Find out more...** about women sexual and reproductive health.

**Sexual Health Info Link: NSW** Sexually Transmitted Infection Programs Unit (STIPU)

Phone: 1800 451 624

Website: www.shil.nsw.gov.au

This website and hotline provide resources, access to information and education and the contact details for services related to STIs.

#### Women and Sexual and Reproductive Health: Position Paper AWHN 2012

This paper advocates for a rights-based approach to ensuring all women can access comprehensive sexual and reproductive health care appropriate to their needs, regardless of their location, age, sexuality, financial status and religious and cultural background.

#### 2.7 Assertive follow-up, outreach and aftercare

Continuity of care when working with women in AOD treatment can take a number of forms, but what is important is the consistent provision of quality care over time. In practice, there are three interventions that support women when they are in the community that provide support whether they are actively involved in other AOD treatment or not. These are:

#### 1. Assertive Follow up

A process of checking in with clients via SMS, phone call, email or letter as a means of support or following up missed appointments.

#### 2. Outreach

A type of AOD service provision that seeks out women who may not otherwise access their service, usually conducted as a mobile service.

#### 3. Aftercare

The provision of support and/or services after or at the conclusion of a specific AOD treatment program, it may take the form of a support group, leisure activity, supported accommodation or case management.

Assertive follow-up for those who miss appointments can help by providing the opportunity to reschedule, to identify and discuss any problems, and to remain engaged. Post-treatment outreach and aftercare services provide the opportunity for transitional support back into the community and to consolidate skills learnt in the treatment setting, including relapse prevention, problem-solving, harm-reduction and self-help skills. Women with complex needs have an increased chance of experiencing setbacks, so post-treatment follow-up can be key to helping them maintain healthy treatment outcomes.

Best practice research indicates that people who participate in follow-up or aftercare after exiting a service have better long-term outcomes. Structured aftercare programs can help provide additional support for people during the transition from treatment into the community. Family, carers or other support people in the person's life should be involved in aftercare planning and treatment where possible (NADA, Complex Needs Capable 2013).

Participation in self-help and aftercare programs is a strongly supported modality for female substance users. Self-help participation was more strongly associated with moving from 'using' to recovery for women than for men in substance use treatment. This finding is consistent with the results of a longitudinal study showing that women seeking help for alcohol problems were more likely than men to participate in self-help groups and to have greater reductions in drinking concurrent with their self-help participation at both 8 and 16-year follow-ups (Grella et al. 2008).

# 3

### IMPROVING ACCESS AND RETENTION FOR WOMEN FROM DIVERSE BACKGROUNDS



- Recognition of diversity and a demonstrated understanding of the issues experienced by women from diverse backgrounds are essential in creating a sense of safety, support and trustworthiness at a worker and organisational level.
- Diversity is a strength that can both motivate and sustain improvements in a client's life and in relation to their general wellbeing.
- Cultural sensitivity and competency are essential to the AOD treatment workforce.
- "Ask the question..." Enquire about potential diversity and show an interest in the response (including expressions of social and cultural identity, sexual orientation and gender identification).
- Working effectively with diversity is about adapting best practice approaches through collaborations with representatives from the specific groups you wish to target.
- Family and community-inclusive practice facilitates improved outcomes for women from diverse backgrounds.
- Forging strong partnerships with specialist diverse population services is an effective way of drawing on already established networks and best practice strategies.
- An awareness and appreciation for diversity among the clients that access AOD treatment can allow for unique opportunities for intervention.

Identifying the qualities that are associated with specific social and cultural backgrounds can become a focus for building identity, belonging and empowerment. Diversity and the related best practice approaches with specific diverse populations of women are explored in this section.

## 3.1 Working effectively with Aboriginal and Torres Strait Islander women

Cultural awareness and sensitivity are central elements for working effectively alongside Aboriginal and Torres Strait Islander women. "Cultural competency can be understood to involve working within a framework that recognises and respects the central importance of culture and identity to Aboriginal and Torres Strait Islander people and communities, working in ways that safeguard the importance of culture, and supports Aboriginal and Torres Strait Islander people's capacity to strengthen the place of culture and identity in promoting social and emotional wellbeing." (NIDAC 2014).

The complexity of issues faced by many Aboriginal people include high rates of trauma (including intergenerational trauma), histories of family separation or disruption and elevated experiences of grief related to early deaths and high rates of incarceration (Lee et al. 2013). "Past Australian government policy and practices enabled the destruction of Indigenous kinship groups, languages and cultural rituals through the forcible removal and separation of Aboriginal and Torres Strait Islander children from their families and kinship networks over some five generations (Ranzijin et al. 2010). The Bringing Them Home report (HEROC 1997) concluded that the forcible removal of children was an act of genocide and that the consequences for Indigenous peoples continue to reverberate today" (NADA 2013). Overt recognition and compassion for the historical experiences of oppression experienced by Aboriginal people by AOD treatment organisations is a useful starting point from which to improve work practices with Aboriginal women.

Indigenous women commonly have many responsibilities within their extended families and household, often having main responsibility for looking after the health of children and other family members (Burns et al 2010). Aboriginal women may also have increased concerns and fears of intervention from child protection agencies and other services, which act as a barrier to their engagement in treatment (Lee et al. 2014, Allan & Campbell 2011).

Data taken from closed drug and alcohol treatment episodes for women in Australia for the period 2012-2013, suggests that Aboriginal women use drug and alcohol treatment services at a higher rate than non-Aboriginal women. This potentially reflects their elevated need for treatment and supports the need to offer tailored programs that are culturally as well as gender appropriate. Aboriginal women have poorer physical health and higher rates of psychological distress than non-Aboriginal women, and are significantly more likely to have children with lower birth weight, be admitted to hospital for assault, and are up to twice as likely to die as a result of illicit drug use (Burns et al. 2010).

The National Indigenous Drug and Alcohol Committee identified key principles for the effective AOD treatment with Aboriginal women. They have been adapted from NADA Working with Diversity in Alcohol and Other Drug Settings (2014) resource and presented here:

#### • Evidence-based and evidenced-informed treatment

Adaptations of evidence-based mainstream interventions that integrate culturally specific practices, including traditional values and reference to spirituality have been shown to be more effective than mainstream services. These elements increase the credibility and relevance of treatment to Aboriginal and Torres Strait Islander people (Terrell, 1993; Anderson, 1992; McCormick, 2000; Brady, 1995b; Gray et al., 2014).

#### • Cultural competency, safety and security

For workers and services, cultural competency can be understood to involve working within a framework that recognises and respects the central importance of culture and identity to Aboriginal and Torres Strait Islander people and communities.

#### • Family and community involvement

When working with Aboriginal and Torres Strait Islander people, involvement of family and community members can be pivotal in achieving best outcomes for an individual (Nagel et al., 2009).

#### Aboriginal and Torres Strait Islander ownership of solutions

Indigenous ownership of solutions is identified as needing to occur, from inception and planning, through to implementation and provision, and then monitoring and evaluation of any solutions.

#### Integrated services and partnerships

Given the complex, multiple needs of people with AOD issues, it is important for specialist AOD treatment services and other services to be well integrated to ensure that people receive all of the services and support they need in a timely fashion and in a way that is easy to access.

Recommendations for responding to Aboriginal women are similar to those recommended for non-Aboriginal women, including addressing co-occurring trauma and mental health issues by "holistically and simultaneously, within an Aboriginal worldview, using a strengths-based approach" (Niccols 2012:331), including linking the support of mothers with their children and cultural components. For example, gender mixing in some services may conflict with cultural perspectives (Taylor et al. 2010).

The Community Reinforcement Approach (CRA) and its adjunct program Community Reinforcement Approach and Family Training (CRAFT) are effective behavioural approaches that use social, family and recreation as positive reinforcers for not drinking alcohol – in contrast to the reinforcing factors that may maintain problematic drinking (Meyers et al. 2011). These interventions are considered to be consistent with the influential role that family and community can play in the lives of Aboriginal and Torres Strait Islander people (Calabria et al. 2013). In a NSW-based study among Indigenous adults, over 90% of participants found that this intervention was considered acceptable to be delivered in their community (Calabria et al. 2013).

For Indigenous women specifically, an evaluation of a 'soft entry' drug and alcohol intervention strategy aimed at Aboriginal Australians in an area of rural NSW found that the approach demonstrated great effect, particularly in regard to increasing the number of Aboriginal women accessing drug and alcohol services in that community (Allan and Kemp 2011). Drug and alcohol counsellors provided access to their services at Aboriginal community events, groups and gatherings. Although no treatment outcomes were measured, the study found the soft entry approach provided services in a way that was acceptable to those receiving them and women were particularly positive.



"Our service has achieved greater engagement with Aboriginal women in our community by facilitating groups and offering services in a lively non-threatening environment and encouraging consumers to bring another community member to share the learning experience and build community capacity."

Our day program idea is attractive for Aboriginal women. They like the idea that they don't have to go away [to rehab]... this gives women anonymity and they can be discreet... they've told me "No one knows I'm in a treatment program – I'm just not available for a few hours a day. I say I'm doing a course."

While it is not always possible to implement significant program changes quickly, the instigation of mutually beneficial partnerships with either Indigenous-managed or Indigenous-focused services can be an extremely powerful place to begin. "It is preferable that all drug and alcohol and mental health services identify and form partnerships with a local Aboriginal health service if they do not have an Aboriginal worker within their team. This helps provide culturally appropriate care." (NSW Health 2009). Developing a Reconciliation Action Plan (RAP) is a concrete way to demonstrate a commitment to reconciliation and support for Aboriginal and Torres Strait Islanders by your organisation. Furthermore, actively seeking to employ Indigenous staff can enhance program delivery to Aboriginal and Torres Strait Islander clients.



Find out more...about working with Aboriginal women.

#### Handbook for Aboriginal Alcohol and Drug Work University Of Sydney 2012 (Lee et al. Ed.s)

This book has been written as a practical tool to use in everyday AOD work, and to be combined with individual skills and local knowledge. It offers a detailed look at alcohol and drug work from clinical, through to prevention, early intervention and harm reduction. This handbook takes into account the complex challenges facing workers when helping clients, their families and, sometimes, whole communities.

#### **Aboriginal-specific Community Reinforcement Approach (CRA) Training Manual NDARC** 2014 (Rose et al)

The Community Reinforcement and Family Training (CRAFT), approach has been used in a number of settings with different population groups. The approach includes two components; the Community Reinforcement Approach (CRA) for working with individuals who are at risk of alcohol related harm and Family Training (CRAFT) for working with their families and loved ones. Evaluations suggest this two-sided approach is more effective for encouraging individuals into alcohol treatment and for assisting family members to feel less depressed, anxious and angry. Most importantly CRAFT appears to fit well with Indigenous views of health and health care that are holistic and dedicated to '... the social, emotional and cultural well-being of the whole community in which each individual is able to achieve their full potential as a human being thereby bringing about the total well-being of their Community'.

#### Implementing a soft entry approach to AOD intervention with Aboriginal women Lyndon Community 2014

This presentation discusses the outcomes of delivering a soft entry approach via an existing community group (Apollo House) in an Aboriginal community in Dubbo. Feedback from a focus group comprised of a variety of community members who have accessed drug and alcohol support in this community will highlight the way day to day practice assesses the suitability and applied evidence of the soft entry approach.

#### NIDAC AOD Treatment Report NIDAC 2014

This paper was developed by the National Indigenous Drug and Alcohol Committee (NIDAC) in response to a misperception that effective AOD treatment is not available for Aboriginal and Torres Strait Islander peoples. The paper aims to allay these misperceptions by outlining who can benefit from receiving treatment. what treatment is known to work; key principles that should guide the application of treatment and what constitutes effective treatment for Aboriginal and Torres Strait Islander people.

#### Working with Diversity in Alcohol and other Drug Settings NADA 2014

The resource has been developed to support non government alcohol and other drug services work with the diversity of clients that access our services and represent the NSW population. It contains examples of best practice.

A supplementary document: **Working with Diversity in Alcohol and other Drug Settings** - **Useful Resources** NADA 2015 also provides a comprehensive list of updated resources of use.

#### **Health Infonet**

A web resource that is a 'one-stop info-shop' for people interested in improving the health of Indigenous Australians.

www.healthinfonet.ecu.edu.au

## 3.2 Increasing supports for culturally and linguistically diverse women

The last census conducted in Australia revealed that over a quarter of the population was born overseas (ABS 2012). Multiculturalism is one of the themes of Australian life however the responses to the needs of people from different cultural heritages have not kept pace. That culturally and linguistically diverse (CALD) communities' have low levels of engagement with AOD treatment services is reflected in national data on treatment utilisation. Although CALD populations may have lower alcohol and other drug use issues when compared to Aboriginal, Torres Strait Islander and Anglo populations, it is apparent from treatment data that those who do experience difficulties are less likely to seek or complete treatment due to cultural attitudes, religious beliefs and conventional patterns of use.

Women from CALD communities commonly experience two key issues related to potential problematic drug or alcohol use: the stress of migration and mental health problems (Browne & Renzaho 2010, Flaherty et al. 2010). Among those in the CALD community, refugees and asylum seekers in Australia are particularly vulnerable to, and affected by, homelessness. Women from CALD backgrounds may also experience violence at increased rates and be at greater risk of victimisation than women from non-minority backgrounds (Roarty & Saggers 2009). Women from CALD backgrounds come into contact with AOD treatment services often as a result of the substance use of a family member or loved one (Lee et al. 2014).

A range of issues have been identified that present barriers for CALD communities accessing treatment, and highlight specific needs and areas relevant for treatment practices. They include:

- language difficulties and lack of availability of translation services
- social stigma
- lack of family and/or community support
- insecure living conditions
- immigration statuses that impact on eligibility for many social and health supports
- cultural perceptions including spirituality and beliefs around health and/or drug and alcohol
  use that may be different to western models of health and harm reduction approaches
  (Roarty & Saggers 2009, Flaherty et al. 2010)
- for older women in particular, language barriers and the social stigma attached to seeking treatment can both be major factors in accessing help.

More time may be needed to build the trust and rapport of CALD women. Also, additional services such as long-term English-language learning and one-on-one case worker support are recommended to assist women to engage in treatment and support their long-term recovery (Roarty & Saggers 2009). Best practice would also indicate the need for training in the area of cultural sensitivity, which enables workers to better appreciate the influence of culture on identity and the experiences of women from different cultural backgrounds.

Many of the strategies and best practice approaches already indicated as effective for women more generally (eg, trauma informed services) are also relevant for services aimed at engaging women from CALD backgrounds. Experiences of trauma specifically related to war and state-sanctioned violence may require additional specialist assistance which may be sought through partnerships with refugee-related services. It is particularly important that treatment programs build on women's strengths, use supportive rather than confrontational approaches, and involve women-only groups or mixed-gender programs that incorporate strong policies related to sexual harassment and safety (Roarty & Saggers 2009). Attention needs to be given to family structures and different cultural approaches to parenting within the context of adapting to the Australian culture. Experiences of difference around the family system and an Australian approach to health and family violence may also be barriers to seeking assistance.



Services in NSW have reported variable success in attracting women from CALD backgrounds. Feedback from services involved in the needs analysis that informed this resource identified a lack of understanding about women's cultural backgrounds and lack of research into the literacy needs of women from CALD backgrounds as gaps in service provision for these women.

We have good mix of staff with different languages... it does help to have a team with different languages.

We try to celebrate the different cultural events as we have women for 12 months...

We don't capture as many [women from CALD backgrounds] as we would like to... we don't recruit well either...

This feedback from the frontline is a good indication of areas needing improvement in order to work effectively with CALD women. It also allows us to put forward some solutions:

ISSUE TO BE ADDRESSED	INTERVENTION
The need to learn and understand different cultural backgrounds and the experience of living in a different culture	<ul> <li>Engage women to 'tell their story' specifically in relation to the story of their culture and what it is like to be in Australia.</li> <li>Celebrate different cultural events, foods, practices, leisure activities, films within the context of the therapy.</li> </ul>
Language barriers	<ul> <li>Examine recruitment processes for staff and volunteers from different cultural backgrounds and with different language skills.</li> <li>Develop resources in different languages.</li> <li>Foster the use of peers from CALD backgrounds.</li> <li>Find out about services with bilingual staff.</li> </ul>
Lack of supports and networks	<ul> <li>Engage specialist social and cultural organisations and ethno-specific agencies.</li> <li>Explore cultural support networks unrelated to AOD treatment.</li> <li>Include CALD training in regular professional development plans.</li> <li>Identify women-only services or gender-separated programs if preferred.</li> </ul>



#### Find out more...about CALD communities

#### Respect: Best practice approaches for working with culturally diverse clients in AOD treatment settings DAMEC 2014

Simple tips for working with clients from culturally and linguistically diverse backgrounds in alcohol and other drug treatment settings. This broadsheet draws on DAMEC's previous research, service evaluations and sector-wide consultation processes to draw together 10 key strategies for practitioners.

Website: www.damec.org.au

Common Threads, Common Practice Best Practice Guide: Working with Immigrant and Refugee Women in Sexual and Reproductive Health Multicultural Centre for Women's Health(MCWH) 2012 (Hach)

Common Threads, Common Practice provides a concise, easy to use reference guide for best practice when working with immigrant and refugee women in sexual and reproductive health. It is informed by the Common Threads Research Report published in 2012 by MCWH, which is also available for download.

Website: www.mcwh.com.au

#### Working with Diversity in Alcohol and other Drug Settings NADA 2014

The resource has been developed to support non government alcohol and other drug services work with the diversity of clients that access our services and represent the NSW population. It contains examples of best practice.

A supplementary document: **Working with Diversity in Alcohol and other Drug Settings - Useful Resources** NADA 2015 also provides a comprehensive list of updated resources of use.

**Working with refugees family project** Australian Centre for Child Protection, University of South Australia 2009

This report outlines the findings of a three stage research project designed to examine why recently arrived families from refugee backgrounds are presenting to the child protection system and to identify culturally appropriate strategies and models for intervention.

**Talking therapies for people with problematic substance use** Te Pou o Te Whakaaro Nui 2010 This guide provides useful background information on problematic substance use, suggestions for interventions during the various stages of treatment, and information on the range of talking therapies currently utilised by practitioners in New Zealand. Also includes sections on Maori, other Pacific Islander cultures and working with refugees.

Helping asylum seeker and refugee background communities with problematic alcohol and other drug use: A guide for community support and AOD workers QNADA 2015 (Pont B and Jaworski A)

The guide is divided into two sections, the first is for workers supporting people from refugee backgrounds across the general health and community services sector who may not have specific training or experience in alcohol and other drugs service provision. And the second is for those working in the AOD sector and discusses issues that are relevant across the range of treatment settings.

## 3.3 Enhanced service-provision for women from LGBTI communities

Research literature that explores substance use among Lesbian, Gay, Bisexual, Transgender and Intersex (LGBTI) communities indicates that lesbian and bisexual women show higher levels of risk in relation to alcohol consumption and problematic drinking than heterosexual women (AIHW 2011; Ritter et al. 2012). Furthermore, according to the Australian Longitudinal Study of Women's Health (ALSWH) lesbian and bisexual women report higher tobacco use, a higher likelihood of having used illicit drugs (41% vs. 10%) and a higher likelihood of ever having injected drugs (11% vs. 1%) when compared to heterosexual women (Hillier et al. 2003; Hughes et al. 2010; Mooney-Somers et al. 2013). It is unsurprising then that the National Drug Strategy 2010-2015 refers to the LGBTI population as a disadvantaged or marginalised group that experiences difficulty accessing drug treatment and achieving successful outcomes from this treatment (Ministerial Council on Drug Strategy 2011).

International research has shown that while LGBTI people may have greater unmet substance use treatment needs than non-LGBTI people, they are often reluctant to seek treatment because of expectations of prejudice and discrimination from health care providers (Couch et al. 2007; Burgess et al. 2008, Grella et al. 2009, Senreich 2009 – source from ACON AOD Strategy). A study of young same sex attracted women and men in Sydney showed that while drug treatment utilisation was high compared to estimates in the general population, it was low relative to the proportion of participants reporting harmful patterns of substance use (Lea et al. 2013).

"Preventing discrimination and stigma is an essential aspect of any comprehensive approach to reducing AOD and MH (Mental Health) problems amongst GLBT. Measures which reduce the stigma and discrimination against GLBT people are likely to have powerful public health impacts." (ACON 2013).

A range of specific characteristics and treatment needs have been identified for LGBTI substance users. Due to the elevated prevalence of drug and alcohol use in the LGBTI community, and potentially earlier age of commencing use, LGBTI clients are likely to enter treatment with more severe substance abuse problems than non-LGBTI clients and have an elevated prevalence of mental health issues (Green & Feinstein 2012). Socialising in the LGBTI communities often occurs in settings where drugs or alcohol are available, increasing women's risk of substance use and relapse after treatment (Senreisch 2009; Drabble & Eliason 2012).

People from LGBTI communities face unique barriers when seeking and staying in substance abuse treatment, including providers who hold stigmatising attitudes about homosexual behaviour or who lack knowledge about the LGBTI community and their health needs (Senreisch 2009; Drabble & Eliason 2012; Green & Feinstein 2012). Furthermore, elevated levels of substance use in the LBTI community can lead to unique expectations and perceived normality around drug and alcohol use, increasing the likelihood of individuals choosing to drink heavily or use drugs and consequently affecting treatment goals.

A comprehensive literature review on the prevalence of mental health issues, AOD problems, and intervention responses in the LGBTI community conducted by NDARC in 2012 concluded that:

- Prevention is a priority principle with LGBTI people; both AOD and mental health (MH) problems are preventable, and interventions such as supportive counselling during adolescence are likely to reduce the risk of later mental health or substance misuse problems.
- Preventing discrimination and stigma is an essential aspect of any comprehensive approach to reducing AOD and MH problems among LGBTI people.
- All AOD and MH services should be LGBTI sensitive. This entails ensuring an adequately trained workforce, culturally appropriate services and a non-judgemental attitude by all staff across the service.
- The variety of treatment interventions, such as (CBT), motivational interviewing,
   12-step programs, and the community reinforcement approach have all been shown to be effective with LGBTI people.

Best practice guidelines to treat LGBTI women with alcohol and drug issues also emphasise the need for cultural competence and inclusive approaches, especially in response to the elevated potential of these clients having histories of trauma and experiences of stigma based on both sexism and homophobia (Eliason et al. 2009). Services are advised to develop trauma informed treatment and consider sexuality and gender issues in assessment procedures and treatment planning (e.g., issues of 'coming out', the role of sexual minority bars, the need for clean and sober social networks for recovery).



#### Case study

Lenny is a 33-year-old single female who identifies as a lesbian. Two months ago she exited a residential rehabilitation program where she had been after an alcohol use relapse. Over the past 10 years, she has accessed residential treatment on several occasions for what she describes as "alcoholism". Lenny was diagnosed with a bipolar condition in her early twenties and regularly sees a psychiatrist for medication reviews. She is in maintenance stage of recovery, attends a 12-step group daily and would like to develop more strategies to assist her in remaining abstinent and also stop smoking.

Lenny is socially isolated from both the LGBTI community and the general community with most of her social relationships being with other 12-step group members. She sees her isolation a consequence of her difference – her shyness, mental health condition and sexuality all making her feel marginalised and she reports that she does not feel "normal". She is conscious of having felt this since early childhood.

Lenny has identified that her harmful drinking increased when she first started frequenting lesbian venues and social scenes as it helped her feel less socially anxious and more confident. The long-term impact of her substance use has added to and reinforced her negative self-beliefs of being incapable and being different. Lenny has expressed a desire to access an LGBTI-inclusive service to maintain her goal of abstinence. Long-term goals for Lenny are to find a meaningful purpose for her life through study and work; establish independence and autonomy from her family while maintaining the strong relationship she has with her parents; and to feel confident enough in her sexuality to pursue intimate relationships with others.

The intervention approach to working with Lenny includes individual counselling (Acceptance and Commitment Therapy) designed to explore her sexuality; a referral to the GP for nicotine replacement therapy and Quit supports and referral to Rainbow Recovery (an LGBTI 12-step group). This combination of treatment interventions provided to Lenny gave her the opportunity to expand her peer support within an LGBTI safe environment place.

The most meaningful progress the client has made continues to occur outside the counselling room. Her skills and abilities in a particular creative field have been recognised by others who have encouraged her to pursue a training and career opportunity in that field. Having a goal that is beyond abstinence and is connected to an endeavour she values has brought meaning and purpose into her daily life.

The case study explored above corresponds with many of the best practice approaches outlined in **Section 2**. While it is apparent that the treatment provided occurred in the context of a LGBTI inclusive service, it is in the exploration and acknowledgement of sexuality that improved practice begins.



#### Find out more....

**ACON** has a range of resources, support services and research projects you can access to improve your service knowledge and service provision regarding the LGBTI Community. Website: www.acon.org.au

#### 3.4 Transgender women

Because gender is assumed to be fixed and fundamental, it can be very challenging to people when a person explores different ways of expressing or experiencing gender, or seeks to change their gender altogether. The term transgender has become an umbrella term to encompass many diverse conditions and identities including describing anyone who lives, has lived, or wants to live, as a member of the opposite gender. A person who identifies their gender as different to what was assumed at birth may consider themselves transgender or trans. An important note for the purposes of supporting clients who engage in AOD treatment is that "everyone has a sex identity and gender expression, regardless of whether they are trans. Everyone has the right to determine their sex identity for themselves, and to express their gender in their own way, without fear of discrimination" (Hyde et al. 2014).

For many people, the terms "gender" and "sex" are used interchangeably, and thus incorrectly. This idea has become so common, particularly in western societies, that it is rarely questioned (gender centre; gender spectrum). Whichever theory people accept as being behind an individual questioning their gender identity, it is important to understand that when a person expresses their identity they are expressing their inherent sense of who they know themselves to be, irrespective of their assigned biological sex.

One of the key issues communicated to the Gender Centre by those transgender women they support is where an AOD service insists that only post-operative transgender women are "real enough" to be entitled to engage with a service as a woman.

"We have had very negative experiences with a number of live-in programs where pre-operative trans women (who have been living as women for over 20 years) have been refused service by a women's AOD program due to their surgical status. What is interesting is that these woman had all of their documentation affirming their female identity, including a passport, and the services still refused access."

Not all transgender people will seek to resolve their gender and sex incongruence with medical intervention. Some people do not experience high degrees of discomfort with their physical bodies and rely on only social transition. Others may not be able to undertake medical transition due to cost or other factors such as other health concerns. In mixed gender services there is a tendency to accommodate transgender women in with males or single rooms where this may not necessarily be a suitable option. Particularly in instances where access to a service is contingent upon availability of a single room, and shared room are readily available.

The Gender Centre outlines these specific tips for enhancing AOD worker support for transgender women:

- Build trust and rapport by being open, sensitive and respectful.
- Don't assume gender or sexuality, ask open questions: "How do you identify your gender?" "what is your current gender identity" (male/female/non-binary/something else\_\_\_\_) followed by "what sex/gender were you assigned at birth?" (male/female), and reflect as an organisation as to whether the second part of this question is essential to the care being provided.
- Ask what pronouns the person would like to be used in the case of a non specified gender.
- Focus on behaviour rather than labels or perceived roles, and remember that all people change and develop along their life cycle.
- Avoid forcing any person to act in ways which are congruent with their birth sex but incongruent with their identity as it will not resolve their feelings of incongruence and may lead to poor outcomes such as depression, anxiety, poor self-esteem, self-harming behaviours and issues of suicidality.
- Offer reassurance and acknowledge the "bravery" of the person in choosing to disclose.
- Assure the person of your confidentiality and make it very clear with whom this information must be shared.
- Allow the transgender person to express all the emotions they are feeling and reassure the person that this experience is not unique they are not alone.
- Ask what support the person has and what supports they feel they are likely to need.
- Indicate your acceptance of the person; acknowledge their diversity and validate it by letting them know it is okay to feel the way they do.

One of the key areas for reform identified by The First Australian National Trans Mental Health Study (2014) was "Government agencies, service providers, and other organisations should ensure that their staff are adequately trained to work with trans people in a respectful and affirmative manner. They must develop policies to actively prevent discrimination and to make trans people feel welcome when accessing their services."



Find out more....about positively engaging with transgender women.

#### The Gender Centre

The Gender Centre offers a wide range of services to transgender and gender diverse people, their partners, families and friends in NSW. They also act as an education, support, training and referral/resource centre to other organisations and service providers. The Gender Centre aim to provide a high quality service which acknowledges human rights and ensures respect and confidentiality.

Website: www.gendercentre.org.au

**The First Australian National Trans Mental Health Study: Summary of Results** Curtin University 2013 (Hyde Z, et al)

This report provides an overview of the findings of The First Australian National Trans Mental Health Study. The study was designed to investigate the mental health and wellbeing of trans people living in Australia.

# 4

# PREGNANCY AND PARENTING IN AOD TREATMENT



- All AOD workers can benefit from having an awareness of health care issues relating to alcohol and drug use in pregnancy.
- Providing factual information that is support-oriented, can encourage women to seek assistance via AOD treatment and other specialist services.
- Women with problematic alcohol and other drug use who disclose pregnancy or child care responsibilities should be provided education, brief intervention and referral to specialist medical assistance with support.
- A coordinated care/case management approach assists with managing the multiple supports that are necessary for providing best practice treatment for women who are pregnant and/or have children in their care.
- Pharmacotherapy is best practice for pregnant women who are opioid dependent.
- There should be a no wrong door approach to accessing treatment.

Women who use alcohol and other drugs while pregnant and/or who have children in their care face stigma and discrimination that can impede AOD intervention. While tackling societal values is an ongoing challenge for women and workers in the AOD sector, there are some fundamental approaches and strategies that can be used to improve outcomes for women, children and their loved ones and this will be explored in this section. Making factual and support-oriented information available to women will educate them about the harms associated with substance use in pregnancy and can show how sustained/problematic use can impact upon parenting.

Fortunately, there is evidence that for some women, pregnancy and parenting can be influential motivators for change in relation to their substance use, entering and engaging with treatment (Greenfield et al. 2007, Mitchell et al. 2008, Jackson & Shannon 2012). In a study of over 100 pregnant women entering inpatient detoxification, qualitative data indicated pregnancy was the top motivator for seeking treatment, with other motivators being recognition of needing help, family, and being tired of the lifestyle (Jackson & Shannon 2012). Similarly, another US study comparing drug using women who were pregnant and not pregnant found pregnant women were more than four times as likely to express a desire or motivation for treatment (Mitchell et al. 2008).

# 4.1 Alcohol and other drug use in pregnancy: Improving pre-natal care outcomes

According to the latest results from the National Strategy Household Survey (2014), over 50% of pregnant women consumed alcohol before they knew they were pregnant and 1 in 4 continued to drink even once they knew they were pregnant. Of those who did consume alcohol, most (96%) usually consumed 1–2 standard drinks per day (AIHW 2014). A small minority of women that continue to drink at high levels are likely to require support to reduce consumption. These women face significant stigma; "Such is the stigma and guilt associated with alcohol use during pregnancy that only a minority of pregnant women with alcohol use disorders access treatment." (Burns & Breen 2013).

Pregnant women who use drugs or alcohol are more likely than non-users to be smokers, to have a psychiatric condition, to be single, Aboriginal, experience greater social and economic disadvantage, to be living in a metropolitan area, younger, unemployed, Australianborn, and to have previously given birth (Kennare et al. 2005; Taylor et al. 2012). Access to a comprehensive range of well-connected services, such as social support, case management, and mental health services for these women is best practice.

Pregnant women with drug and alcohol issues require services that can support them throughout their pregnancy and provide post-natal support for themselves and their baby. Among pregnant women who use drugs or alcohol, opiate use remains most prevalent; however, use of other drugs, such as methamphetamine, is of increasing concern due to their impact on women's mental and physical health and the heightened risk they pose for obstetric complications (Taylor et al 2012).

For those organisations that do not have the capacity to cater for women who use substances and who are pregnant or who have children in their care, there is still the opportunity for brief intervention, education and supported referral. The specific and complex needs of pregnant women who use alcohol and other drugs requires a coordinated response using a multidisciplinary team approach to include specialists across neonatal, drug and alcohol, mental health and social work fields. Applying a coordinated case management or care management approach can assist in bringing together the specialist services needed by women that the AOD treatment service cannot provide in isolation. Care coordination involves deliberately organising client care activities and sharing information among all of the participants concerned with achieving safer and more effective care. This means that the client's needs and preferences are known ahead of time and communicated at the right time to the right people, and that this information is used to provide safe, appropriate and effective care to the client.

Women on pharmacotherapy need information that empowers and supports them to make choices and decisions about how their pregnancy, birth and the period after birth is managed. Regular benzodiazepine use during pregnancy is associated with neonatal abstinence syndrome as are opioids. However, benzodiazepines may be administered for symptomatic relief during inpatient alcohol withdrawal if required with appropriate supervision and neonatal care (NSW Health 2015). For pregnant women who are dependent on benzodiazepines, best treatment approaches aim for tapering of dose and subsequent discontinuation of use if and when this is safely achievable. Benzodiazepines are generally unsafe for use while breastfeeding and can cause significant withdrawal symptoms for the newborns of dependent mothers (Keegan et al. 2010).

For opioid-dependent pregnant women, induction to methadone maintenance as part of comprehensive treatment programs is considered best practice (Minozzi et al. 2008; Keegan et al. 2010). Commencement of methadone should ideally take place in an outpatient setting, with maternal and obstetric supervision - as is common practice in Australia, and supported by the American College of Obstetricians and Gynaecologists (ACOG). Continued buprenorphine treatment is safe for women who become pregnant while already in this type of treatment, achieves similar outcomes in reducing maternal drug use and promotes positive neonatal outcomes to methadone, and appears to result in a less severe neonatal abstinence syndrome compared to methadone (Jones et al. 2012; Young & Martin 2012). However, as buprenorphine induction may place the foetus at risk from precipitated maternal withdrawal and potentially trigger relapse in mothers, methadone is recommended for active opioid users who need to be inducted to pharmacotherapy during pregnancy (Young & Martin 2012).



#### Case study

Sally is 19 years old. She first presented to the service when she was 17 years old with a long history of substance use – including cannabis use from age 13, heroin use at age 15 and misuse of prescribed benzodiazepines. Her eldest sister died of a heroin overdose when Sally was 15 and her parents separated. Sally was engaged in a methadone program at the time she presented for treatment. She also reported experiencing multiple mental health issues including panic attacks and periods of feeling suicidal.

While Sally's continued substance use has had an impact on her family relationships in the past she has a supportive relationship with her mother, younger sister and her mother's partner. Support from her stepfather has included providing childcare for Sally's young daughter while she attends a support program and, more recently, TAFE.

In spite of initial lack of engagement with the AOD treatment service, assertive follow-up techniques were employed, flexibility in terms of accessibility was provided as well as a harm minimisation approach. Counselling appointments were arranged in order to develop strategies to manage her high anxiety, creating trust with the service and support to stabilise her drug use so that she could attend a day treatment program.

On commencing the program Sally discovered she was pregnant, with no support reported from the father. Sally's pharmacotherapy clinic referred her to the Chemical Use in Pregnancy Service (CUPS) which has links with FaCS. She was required to have contact with them throughout the pregnancy and she was advised that Family and Community Services could become involved if there was any indication of drug use throughout the pregnancy and concerns once the baby was born. This then led to a series of monitoring mechanisms being put in place with the expectation that Sally comply. This included weekly appointments to the CUPS service and random drug screen urine testing.

Sally responded to the news of the pregnancy by immediately stopping all use of illicit substances. She committed to maintain stability on the methadone treatment and to only take the benzodiazepines as prescribed. She also began the day program. Her mother became involved in supporting Sally to meet the obligations of her ante-natal care plan. The progressing pregnancy and her experiences with the ante-natal system resulted in Sally requesting extra support. This support involved such things as securing practical equipment and needs for the expected baby as well as support and advocacy in navigating the hospital and ante-natal systems. It also included providing information relevant to her about pregnant women on methadone programs.

Sally completed the day program and then participated in an aftercare group until the birth of her child. Having the ongoing support of AOD workers throughout the process assisted Sally in coping with experiences of stigma and discrimination during the birth and in relation to breastfeeding. Sally continued to attend aftercare programs and brought the baby to groups.

This case study highlights the importance of working with the client where they are at. For Sally to be able to establish an effective therapeutic relationship with an AOD worker, they needed to be flexible, patient and yet persistent in the early stages of engagement. The role of advocacy became vital to ensuring that Sally receive access to the ante-natal care that any pregnant woman is entitled to. When this was denied, the role of the worker was to challenge and ask why.

The outcomes of pharmacotherapy interventions are significantly enhanced for pregnant women with the support of evidence-based psychosocial interventions and a range of other supportive processes including:

- treatment engagement strategies
- treatment approaches that begin during pregnancy
- programs that incorporate standard evidence-based practice treatment modalities
- home-based interventions
- comprehensive treatment that includes perinatal care, mental health services, child-care and other supports
- service integration approaches, where addiction services are provided with onsite services related to pregnancy, parenting, or other child-related services.

At the time of writing, there are a number of specialist services for pregnant women who use drugs and alcohol in NSW, including those that cater for pregnant women enrolled in an opioid pharmacotherapy program. Specialist services include:

#### • Chemical Use in Pregnancy Services (CUPS)

Langton Centre and Royal Hospital for Women - Sydney (CUPS: (02) 9382 7111, Langton Centre: (02) 9332 8777, Caringbah (02) 9540 7668)

#### Western Sydney and Westmead Hospital Drug Use in Pregnancy Service (DUPS)

Westmead and Blacktown

(DUPS: (02) 8860 2560 or (02) 8860 2565)

#### Substance Use in Pregnancy and Parenting (SUPPS)

Wollongong

(SUPPS: 1300 652 226)

#### • Perinatal and Family Drug Health (PAFDH)

Royal Prince Alfred Hospital (PAFDH: (02) 9515 7611)

#### South West Sydney LHD Perinatal and Family Drug Health (PAFDH)

(PAFDH: (02) 9616 8586)

#### • Mid North Coast LHD Drug Use in Pregnancy Services (DUPS)

Kempsey 1300 662 263 Coffs Harbour 1300 662 263 Port Macquarie 1300 662 263

#### Northern LHD

Tweed Heads Community Health Mum's Program (07) 5506 7823 Riverlands Detox Centre Lismore (07) 6620 7608

Gold standard treatments for women who are pregnant and use substances include extended hospitalisation if required, additional support during birth and post-delivery, and the option of assertive follow-up of the mother and child through the child's formative years to assist the woman in the areas of healthcare, navigating social services, accessing housing and positive parenting (Burns & Breen 2013). Women-only and gender-specific treatment programs are best suited to respond to the specific health and medical needs of pregnant women with substance use issues (Greenfield et al. 2007).

The Clinical Guidelines for the Management of Substance Use During Pregnancy, Birth and the Postnatal Period, released in 2014 is the key resource in NSW when working with AOD use in pregnancy. The guidelines have been developed based on "the best currently available evidence, developed through a rigorous process in which international and Australian research literature was reviewed by experts and consensus achieved."



**Find out more...**about substance misuse in pregnancy.

#### Clinical Guidelines for the Management of Substance Use During Pregnancy, Birth and the Postnatal Period. NSW Ministry of Health 2014

These guidelines contain information on the use of buprenorphine maintenance treatment during pregnancy; cannabis use in pregnancy; a recommendation for abstinence from alcohol during pregnancy; and updated information on child protection legislation in NSW. New sections include prescription opioids use in pregnancy; use of antidepressants in pregnancy; management of withdrawal; management of incarcerated pregnant women; and management of women in rural and remote areas.

#### It's time to have the conversation: Understanding the treatment needs of women who are pregnant and alcohol dependent. (NDARC 2013).

This report presents a narrative literature review of treatments available to pregnant women who have alcohol use disorders and findings from interviews with key stakeholders regarding current treatment practices and areas requiring improvement.

#### Supporting Pregnant Women who Use Alcohol or other Drugs: A guide for primary health professionals NDARC 2014.

This guide is intended for a range of health professionals, in a variety of settings to help support and provide information to pregnant women who use alcohol and other drugs. It also contains a detailed resource list of additional support services and resources.

**MotherSafe** is a free telephone counselling and advice service for pregnant and breastfeeding women (9am-5pm) based at the Royal Hospital for Women, Randwick.

The MotherSafe website also hosts a number of factsheets for alcohol and other drug use in pregnancy – Alcohol, Cannabis, Tobacco, Opioids, Amphetamines, Benzodiazepines.

Phone: 9382 6539 (Sydney Metropolitan Area); 1800 647 848 (Non-Metropolitan Area); Non-English speakers can access an interpreter through TIS National by phoning 131 450. Website: www.mothersafe.org.au

# 4.2 Introducing child-sensitive service provision and support for positive parenting

Women who have children and experience issues with their drug and/or alcohol use are more likely to have a range of material and personal support needs; a higher prevalence of psychiatric symptoms; early childhood and lifetime trauma experiences; and lack social support. These issues present significant clinical challenges, and many of these factors may also inhibit the mother's attachment to her infant, the advancement of their parenting skills and their capacity to maintain child custody. In addition, mothers may face difficulties maintaining engagement with, or completing, AOD treatment (Greenfield et al. 2007).

Significant maternal drug or alcohol use is associated with a variety of caregiving, child and family functioning problems, including a greater likelihood of neglect or abuse of children, reduced emotional involvement and attachment, increased punitive behaviour toward children, insensitive and interfering behaviour, ambivalent feelings about retaining custody, feelings of guilt and increased parenting stress (Fraser et al. 2010; Suchman et al. 2011). The continuing presence of drugs or alcohol significantly reduces an individual's dopaminergic response to stress, leaving the mother highly vulnerable to negative emotions and potentially lacking feelings of pleasure or reward ordinarily associated with caring for young children (Suchman et al. 2011).

The need to enhance capacity for care giving, and interventions focusing on the relationship between mother and infant, is a developing area of research. Best practice treatment services for women with children include:

- women-centred treatment that involves children such as women-specific outpatient clinics and day programs; women-only residential treatment including residential services that allow children to stay with their mothers and/or family residential services
- specialised health and mental health services, particularly pre and post-natal health interventions and specialist mental health interventions such as individual or group therapies
- home visits, typically by a nurse, focusing on providing maternal support, promoting healthy parent-child interactions, and providing information and linkages to material resources
- concrete practical assistance, such as transportation, childcare, and worker assistance to link with treatment services
- short-term targeted interventions, including psycho-educational groups, counselling or support groups and contingency management approaches
- comprehensive and holistic interventions, including programs that integrate several of these components.

The following is a summary of a specific parenting program delivered within an AOD treatment setting. The descriptions are sourced from Gruenert and Tsantefski (2012) Responding to the needs of children and parents in families experiencing alcohol and other drug problems.

#### PARENTS UNDER PRESSURE (PUP)

The Parents Under Pressure (PUP) program, developed by Professor Sharon Dawe (Griffith University) and Dr Paul Harnett (University of Queensland), is designed for families with multiple problems and where children are at risk of adverse outcomes. It is an intensive multicomponent program comprising 10 modules, delivered in 12 sessions of 1.5 hours duration each. PUP has been used where there is problematic parental substance use and there are concerns about child wellbeing and protection. The program is delivered by a trained staff member, generally in the family home, and focuses on a range of parental challenges. Beginning with parents' perception of themselves, PUP encourages parents to acknowledge their own strengths and identify and comment on their children's positive behaviours. The program helps parents to identify the various factors that make parenting difficult and devise strategies to overcome these challenges.

This technique differs from the traditional parenting deficit model in that it empowers parents to discover and use their strengths to build positive parent—child relationships, without focusing on a lack of specific parenting skills. PUP includes daily child-focused playtimes and helps parents to improve their mood and use non-punitive methods to control children's behaviour. In addition, the program aims to help families re-connect with their local community, extend social networks and cope with practical family issues, such as child care, school involvement and other social support. The PUP program has demonstrated significant reductions in family stress, child abuse potential, problems with others, and improvements in parent and child functioning and parent—child relationships (Dawes et al. 2008).

Website: www.pupprogram.net.au



**Find out more...**about parenting support in AOD treatment.

The Parenting Support Toolkit is a resource to help alcohol and other drug workers address the vital role that parenting plays in their clients' lives.

- Parenting Support Toolkit: Overview (VIC Health 2011)
- Parenting Support Toolkit: Quick reference card (VIC Health 2011)
- Parenting Support Toolkit: Booklet 1 (VIC Health 2011)
- Parenting Support Toolkit: Booklet 2 (VIC Health 2011)
- Parenting Support Toolkit: Booklet 3 (VIC Health 2011)

**Strengthening Aboriginal family functioning: What works and Why?** Child Family Community Australia, Australian Institue if Family Studies 2014 (Lohoar, Bureta & Kennedy)

This paper makes extensive use of state-specific research and data. There is a particular focus on recent data describing the level of functioning of Aboriginal families in Western Australia.

#### 4.3 Working effectively with Family and Community Services

While it is essential to avoid any assumption that parental AOD use automatically increases risks for dependent children, the links between maternal substance use and the risk of neglect and harm to their children have been well-documented. Parental drug use is considered to be a key factor in recent child death reviews (NSW Child Death Review Team, 2012, 2013, 2014). Other studies into the links between parental substance use and child abuse and neglect have indicated that:

- Parents who misuse substances tend to have poorer parenting styles. Substance abusing mothers are more likely to be more harsh, punitive and inconsistent in their parenting styles.
- Families involved with significant drug use are more likely to be facing unemployment, poverty, housing instability and social isolation.
- Children's medical, nutritional, social and emotional needs are more commonly neglected.
- Family violence is more common in households where there is significant use of alcohol and other drugs. (NSW Community Services, 2010)

The above findings are just a sample of the concerns that can occur where parental drug use impacts on dependent children. It is also important to note that maternal substance use is only one of a number of factors that can contribute to risks associated with neglect and harm of children; factors such as mental illness, history of trauma, socio-economic disadvantage and isolation can commonly co-occur with substance misuse. It is, of course, important to view substance use and any impact this may have on parenting capacity within context rather than assuming that AOD use is the causal factor (Dawe, Harnett and Frye, 2008).

Given the increasing evidence of the potential impact of substance use on parenting capacity and the subsequent risks for children, it is imperative that child protection and family support services across both government and non government sectors be involved in providing support for parents and protection for children deemed to be at risk of neglect and harm.

It is essential that this section begins with the premise that Family and Community Services (FaCS) involvement with substance using women with dependent children is, ideally, a positive protective factor and can lead to significant change and positive outcomes for families.

"I never felt like Kait and Charlotte were just my caseworkers, it felt like we were a team – that we all had the same goal. This made it easier to give up drugs. Kait and Charlotte were looking out for me and I didn't want to let them down." (Hannah, mum in Shining a Light on Good Practice, NSW Family and Community Services, 2014.)

Women with dependent children who are involved with significant levels of substance use are commonly involved with FaCS. It is estimated that among the 309,600 children reported to FaCS in 2008-2009, the prevalence of substance abuse problems amongst their parents ranged between 50 and 80%. A review of the case files families reported to FaCS indicate that in 41% of cases at least one parent was known to misuse drugs, and 46% misused alcohol (NSW Community Services, 2010).

FaCS involvement with children of substance using mothers invariably brings legal pressure on the parent to address their AOD concerns. The Child Protection Amendment Act 2014 increased the directives for parents with dependent children to address factors, such as significant AOD use, deemed to be impacting on parenting capacity. The Child Protection Amendment Act 2014 increased the focus on earlier intervention, expanded Parental Responsibility Contracts to expectant mothers, introduced new court-issued Parental Capacity orders and introduced changes to the permanent placement principles.

Due to a number of factors, not least the inherently coercive role of FaCS in directing mothers to address their AOD problems, there is often a significant level of stress and uncertainty where FaCS are involved.

PRINCIPLES OF STRENGTHENING A WORKING ALLIANCE WITH FACS It is essential for AOD workers to remember that from the perspective of FaCS workers, child safety and well-being is always the key focus. While AOD workers may have the opportunity to work with children, it is more common for AOD workers to build a working relationship with the mother. However workers within both AOD and FaCS services ultimately hold the hope of the best outcome for families and working collaboratively, including resolving disputes related to worker approach through professional channels will support this aim.

In order for AOD and FaCS services to work effectively towards a common goal it is important to reduce the likelihood of combative situations and enhance the working alliance between services by:

- Continuously building relationships at a local level
- Being informed and informing regarding the scope and limitations of services
- Being willing and available to share expertise as required
- Having up-to-date knowledge of relevant legislation, policies and procedures
- Clear records of interactions between services
- Addressing any disputes between services in a professional and appropriate manner.

In situations where issues have been raised with the caseworker and/or the appropriate managers and differences have not been resolved on a local level between service providers, it may be appropriate to make contact with the Manager of the Clinical Issues Team in FaCS (see overpage) to assist in identifying the next steps.

#### **FACS INITIATIVES**

There has recently been a shift in the culture of FaCS at an organisational level under the banner of the "Care and Protection Practice Framework", launched in December 2012. The aim of the Framework is essentially to enhance the skills and knowledge of front-line FaCS Caseworkers and to increase the level of collaboration between FaCS and other services. Embedded within the Framework are ten practice standards that have been developed with the FaCS Office of the Senior Practitioner:

- Standard 1. Practice leadership
- Standard 2. Relationship-based practice
- Standard 3. Holistic assessment and family work
- Standard 4. Collaboration
- Standard 5. Critical reflection
- Standard 6. Culturally responsive practice with Aboriginal communities
- **Standard 7.** Culturally responsive practice with diverse communities
- Standard 8. Practice expertise
- Standard 9. Sharing risk
- Standard 10. Documentation in casework.



The organisations consulted in the development of this resource made a number of suggestions for improving collaboration between AOD services and case workers from FaCS. These included the development of a continuum of care model for women in AOD treatment who are involved with FaCS; education of FaCS workers about best practice in drug and alcohol treatment; increased opportunities for networking and joint training for drug and alcohol workers and FaCS case workers; and a dedicated alcohol and other drugs worker position in FaCS (eg, seconded from the specialist women's AOD service sector) to undertake drug and alcohol assessments and recommend a suitable care plan.

The increasing number of women with children not in their own care (ie, children subject to Emergency Care and Protection orders or long-term care orders) was consistently raised as a key issue in providing drug and alcohol treatment to women. Services also reported that more children are not being restored to mothers, or there is a lengthy restoration period – in some cases taking up to two years for a review of the care order – resulting in significant feelings of grief and loss, which can reduce women's motivation to continue AOD treatment. Services also reported that some women clients become disgruntled with AOD treatment services when children are not restored, which undermines the therapeutic alliance.

A number of services also described cases in which women were unaware of what FaCS required for child restoration, or indeed if restoration was possible, and women were entering treatment of their own accord in the hope that their children would be restored or access visits would be increased.

For four years pretty consistently we have had children vacancies... we haven't been full [with children] because FaCS are removing children... we don't have a child in residency at the moment. We don't get children because they are either on permanent guardianship orders or the restoration looks very slim and they [women] are coming to us in the hope of [restoration].

We are the same. Sometimes we think "we're a child service and we don't have children".

The variable working relationships with FaCS case workers was also raised as an important issue. Some services highlighted a lack of clarity in FaCS case planning for women and their children or a lack of communication to the services about the plan, leaving women and the workers who care for them unsure of the goals set by FaCS and what was required to meet those goals. Some services found it difficult to ensure the presence of FaCS case workers at the service's care planning meetings, while others had established routines with some FaCS case workers from local offices.

They [FaCS] used to turn up with a 17-page care plan and now you never see one. Women have the impression if they do a residential program they will get their kids back and it's not the case today.

Some [FaCS] workers will be engaged, will have a plan with our clients when she leaves us but ... this is at an individual level it is not system wide – we don't have an agreement with FaCS across the state although it'd be great to have that.

At the beginning of a woman's stay we set up an 'expectation meeting' between FaCS, our staff and the client. We ask the case worker what they expect... it's a transparent process and everyone knows the plan. The case worker is called back four weeks before the woman's discharge and a detailed plan is made then too.

The impact of Care and Protection Orders on the children of substance-using mothers was also raised as an important area for consideration by services. Children who expect, but fail, to be reunited with their mothers, and children in care who must travel long distances to see their mothers for access visits were two groups that were highlighted as needing long-term support.

... Continued over page



The reluctance of women to disclose substance use during pregnancy or to seek treatment out of fear of scrutiny by FaCS was also highlighted as a key issue, particularly when it results in women not receiving adequate antenatal care or late detection of neonates in withdrawal.

"We had a woman referred to the residential program for a telephone assessment. It wasn't until a couple of more phone support contacts that she revealed she was pregnant and due in 5 weeks. During the entirety of her pregnancy she was engaged in opiate treatment and attending a methadone clinic daily. For fear of having her baby removed, she had successfully concealed her pregnancy with baggy clothing and had consequently attended no prenatal assessment nor treatment. It's a terrible reality that for this mother to be, the fear of having her baby removed could have had serious implications for the health and wellbeing of the baby".

#### **Practice First**

A critical component of the Care and Protection Practice Framework has involved the roll out of an initiative referred to as Practice First within frontline FaCS services across NSW. The impetus for the Practice First approach was the concern that FaCS case workers needed to spend more time involved with working with families and be less involved with red tape and paperwork. Its focus is on changing the practice culture across the spectrum of work with families including assessment, intervention and collaboration with partner agencies. Practice First aims to achieve safety for children and families through skilful child protection practice, shared management of risk and genuine relationships with families and community.

There are three essential components to Practice First:

- 1. Principles of practice guide the model. They are evidenced based, reflect contemporary research about what works in child protection and provide a solid framework for improved outcomes.
- 2. People and practice leadership casework is delivered by teams, not individuals, and teams have the support of administrative staff and timely clinical and specialist support. Skill development is ongoing and requires practitioners to have insight into the impact of their own practice on families, and to strengthen their skills in working with families to change
- **3.** Realignment of systems a clear mandate giving legitimacy to family work has been emphasised, freeing casework time from administration, and sharing risk and decision making across teams.

Group Supervision is at the heart of Practice First. Caseworkers, managers, casework specialists, psychologists and other specialist staff such as the Clinical Issues team consultants meet each week to discuss cases and reflect on practice. Partner agencies are regularly invited to group supervision sessions, which can include local drug and alcohol services. This genuine partnership has many benefits: it creates more efficient and effective referrals and information sharing, shared understanding of the risks and protective factors, greater transparency around decisions, more purposeful case plans and wrap-around services, and ultimately, better outcomes for children and families.

For AOD workers involved with women and their children, the implications of the Practice Standards and the Practice First initiative should be that there is an improvement in the collaborative nature of the relationship between services and a greater level of transparency in regards to decisions made within FaCS. Drug and alcohol workers should contact the Community Services Centres they are working with to find out how the practice standards might help them in working together. If they are a Practice First site this could include being involved in group supervision when discussing shared clients.

For more information on the practice standards see the **Practice Standards Information Sheet** hosted on the NADA website women's focus area page and read the Creating Better Outcomes article in the **NADA Advocate 2015 June edition**.

#### **Clinical Issues Team**

Within the FaCS Office of the Senior Practitioner there is a specialist unit, referred to as the Clinical Issues Team (formerly the Drug and Alcohol Expertise Unit). The focus of the Clinical Issues Team is to educate and support front-line FaCS workers specifically in relation to working with families where there is significant AOD, mental health issues and/or domestic violence. The team consists of five clinical consultants and a manager who have all had significant clinical experience in AOD, mental health and/or domestic violence outside of the FaCS context. They offer consultation to caseworkers and encourage collaboration between caseworkers and other service providers, as well as attending and group supervision. The team also run training sessions for caseworkers on the clinical issues in Community Services Centres across NSW and encourage caseworkers to invite local service providers to also attend these sessions.



Find out more...about the FaCS Care and Protection Practice Framework

Care and Protection: Practice Standards Information for families working with Family and Community Services caseworkers Office of the Senior Practitioner, Family and Community Services, Community Services NSW

This flyer provides a brief overview of what families working with FaCS can expect from case workers. A new version of this flyer is under development for non government service providers and will be available soon.

Website: www.nada.org.au/nada-focus-areas/women to download the flyer from the NADA website or visit www.community.nsw.gov.au

# 5

# BUILDING STRONG PARTNERSHIPS TO INCREASE POSITIVE OUTCOMES FOR WOMEN



- Women need to be invited to take an active rather than passive role in the development of care plans and in the selection of external support services.
- Establishing strong respectful partnerships with other services can fill the gaps that may not be covered by AOD treatment.
- There are a number of important factors that can support good partnerships such as clear goals, clearly defined roles and responsibilities, and documentation that explains the nature of the partnership and that relates to governance.
- Creating seamless service provision to women can be supported by collaborative care planning and the implementation of supported referral processes.

Women accessing AOD treatment present with multiple complex issues and as a result may be overwhelmed by how they can navigate various services in order to improve their wellbeing and have their needs met. In parallel, many AOD workers also feel confused by the best ways to assist their clients – particularly if they feel unable to provide the services that are required. The most effective and useful response in this situation is to establish strong respectful partnerships with other services that can fill the gaps that may not be covered by AOD treatment such as childcare support, specialised mental health support, practical welfare interventions and health care.

In this section of the resource different types of partnerships are explored. It begins with the most valuable partnership – that formed with women who engage with a service for AOD treatment – then goes on to present specific community-based partnerships, followed by those partnerships that can help ensure effective aftercare.

#### 5.1 Involving women in effective case management

In keeping with a strengths-based approach and the goal of self-determination by women engaged in women's AOD treatment, involvement in case management is essential. What this means in practice is women take an active rather than passive role in the development of their care plans and the selection of external support services.

Women who are central to decision making about their care and treatment are more likely to have an investment in its outcome. Feeling that they have been genuinely consulted and presented with options from which they can make a choice is empowering – and may have a significant impact on the experience of AOD treatment. Some useful considerations for effective case management that actively considers the needs of clients is outlined below by Gelb & Rutman 2011:

- Establish a trusting relationship with the person.
- Establish close and frequent communication with them (e.g. check-ins and regularly scheduled meetings).

- Develop a care plan based on their goals, strengths and needs.
- Facilitate transportation for important appointments.
- Liaise with other service providers with consent.
- Assist the development of life skills and/or the person's participation in healthy, safe activities.
- Establish close communication with supportive significant others in the person's life.
- Advocate for improved communication among service providers, continuity of care and access to care/services.



#### Case study

Samantha is a 27-year-old woman with a history of alcohol use. She has previously engaged in residential treatment, but was discharged for non-compliance and aggressive behaviour towards other residents. Samantha was encouraged by her parole officer to try residential treatment again as she has a number of outstanding criminal justice issues, including a breach of an Apprehended Violence Order against her mother and grandfather. Samantha is concerned about her relationship with her family, particularly her grandfather who she has always been close to but who she recently assaulted when under the influence of alcohol and for which she was charged.

Prior to attending for treatment, Samantha was drinking 10-15 cans of beer every day. She also smoked methamphetamine, but her primary drug of concern is alcohol. She began drinking at age 14 and smoking methamphetamine at age 22. Samantha has engaged well with the support provided to her, including that provided by another resident.

Samantha has body image issues. When she attended the service previously she would wear large clothes to cover her body up and was not eating at all. This time she wore normal clothes, was eating well and focusing on her physical health as one of her goals. Samantha has been supported to identify her immediate treatment goals, which include a plan to rebuild and reconnect with her grandfather. Her other treatment goals include sharing her story with the treatment community and focusing on her physical health.

Samantha has agreed to be referred to a specialist that works with clients with complex needs and that will work with her while she is in treatment. She will also attend appointments with the visiting mental health nurse and the visiting psychiatrist.

In the case study explored above, there is clear evidence of client-led case management. Samantha identified the goals she wanted to focus on and, with support, has started to address them. Also her more complex issues – such as the mental health concerns identified – are more likely to be considered by the client to require intervention if it has been established that it is her decision. For women who have had previous poor experience of AOD treatment, coupled with criminal justice or custodial experiences, it may take time for trust to be built and for them to arrive at a decision to tackle complex issues such as mental health. The benefit of working in this way is that the outcomes are likely to be sustained as the client has made an investment of their own free will.



Find out more...about client-led case management.

#### Specialist Homelessness Services Case Management Resource Kit (NSW Dept. FaCS 2012)

Case management has clear benefits for clients and service providers. It means clients have better access to services and are empowered to make decisions about the support they receive. For service providers, case management reinforces the shared responsibilities of clients and the services that work with them, and helps to identify the client's strengths. The kit aims to provide caseworkers and casework supervisors across all SHS with knowledge and tools to work effectively with people who are homeless or at risk of homelessness.

**Counselling guidelines: Alcohol and other drug issues** 3rd Edition WA Drug and Alcohol Authority (Marsh et al. 2013)

These guidelines includes over 50 chapters exploring some of the key skills needed to work at an individual level with people who have AOD problems including a case management, group work, referral, working with intoxicated and incarcerated clients, clinical supervision and stress and burnout.

#### 5.2 Building community-based partnerships

While it may be possible for an AOD treatment service to put in place changes that reflect gender-responsive, family-inclusive and trauma informed practices, it is not the expectation that AOD treatment has the capacity to address all needs. There are some useful tips in guiding the management of good partnerships across organisations. These tips, adapted from Hunter and Perkins (2014) include:

- clear goals and purpose
- clearly defined roles and responsibilities
- documentation that explains the nature of the partnership
- robust monitoring and evaluation
- sharing of information with clear protocols
- use of 'local champions' within partner organisations
- for long-term partnerships, a clear purpose and governance structure is recommended.

Useful partnerships to consider might be with child/youth organisations that provide support around the substance use of a parent, collaborations with outreach or drop-in services that can provide after-hours support for non-residential services, coordination with women's specialist antenatal care services or other local primary health centres.

"While partnerships can be established at a personal level, sustainable partnerships are likely to require formalisation at the agency–agency level where possible, with clear aims and objectives, inter-agency protocols and information sharing, supported by policies, procedures, and documented referral pathways. It is acknowledged that good partnerships take time and effort to develop" (Grenert & Tsantefski, 2012:10).

Other ways of creating seamless service provision for women is through collaborative care planning (as outlined in **Section 2.5** of this resource) and the implementation of supported referral processes. A 'warm' or 'supported' referral means that there is some personal contact from the worker who is making the referral. This might take the form of an introductory phone call to the service in order to make introductions on behalf of the client, or attending the initial appointment with the client as a support, or it might be that a representative from the service being referred to attends the AOD treatment service to establish some rapport with the client.



We have got to have women's and children's services, but we also need to be looking at better access to childcare so women can attend a day program if they choose.

Working on partnerships that will continue to support women into aftercare (see **Section 2.7**) is essential to sustaining AOD treatment outcomes. For more information and resources related to building partnerships explore the Partnerships information on the NADA website: www.nada.org.au/whatwedo/partnerships.

There is evidence to suggest that women appear more likely to seek treatment in non-speciality settings (Ettorre 2004; Greenfield 2010). This suggests a need for collaboration and/or co-location of drug and alcohol treatment services with other social and health agencies. In turn, these services could offer treatment that address these needs and issues common to women with problematic substance use. Recognising complementary services for women in the community such as GPs, playgroups, childcare facilities and creating support links that encourage referrals to specialist AOD treatment services has the potential to increase accessibility for women experiencing substance use issues.



Find out more...about building strong partnerships.

#### Promoting partnerships: Mental Health Coordinating Council

MHCC is working towards promoting partnerships in two ways:

- Building structures and processes that enhance effective partnership behaviour.
- Building capacity in the sector to work in effective partnership arrangements.

This section of their website explores in detail these two aims with supplementary resources for each stage.

Website: www.mhcc.org.au

#### No Wrong Door (Ovens & King Community Health Service, Victoria)

The No Wrong Door website has various functions, with its primary focus to enable workforce capacity to better identify, assess, respond and treat people with dual diagnosis presentations. The website provides a platform to host regional policy, procedures, protocols, pathways, committees and educational opportunities for the member agencies. In addition, this website is designed to host a discrete area for consumer and carer representatives to better support them in their roles with their employers.

Website: www.nowrongdoor.org.au

#### 5.3 Identifying partnerships that support effective aftercare

Making the transition from AOD treatment to the community requires planning and a focus on building sustainable community-based supports that will assist women to maintain the changes they have made to their lives. Best practice aftercare planning occurs as part of treatment, specifically in relation to identifying the types of supports that may have been lacking in a client's life prior to entering treatment, such as:

- stable accommodation and/or financial supports
- ongoing psychiatric or psychological support
- general health supports
- vocation, education and/or employment
- childcare supports
- engagement in leisure activities.



I've had several clients who did well over six months [in the program] but they finish and go back to the community, back to their old environments, and there's not enough support... even for clients who are having counselling it's only one hour a week – it's not enough support, and they often relapse.

The biggest problem for aftercare is housing. The Department of Housing policy is to retain the accommodation for three months only and we have a six-month program – if a woman has to give up her property she'll be deterred from coming in – she'll be homeless... If a woman has lost her baby she is less likely to come into treatment for fear of losing her housing, and if they do come in and lose their accommodation, six months isn't enough time to find other accommodation for them.

Identifying potential partnerships that may assist you and your organisation to facilitate these different supports within an aftercare plan, are likely to ensure sustainable treatment outcomes and prevent potential relapse.



Find out more...about building strong partnerships.

#### The NADA website **Partnership Webpage**

Provides information, resources and tools to establish effective partnerships.

Website: www.nada.org.au/whatwedo/partnerships

Joining Together: Tackling AOD harms through a regional partnership (Dunne 2013) Is a case example of an effective regional partnership and the journey taken to get there.

#### The Partnering Toolbook

The Partnering Toolbook builds on the experience of those who have been at the forefront of innovative partnerships and offers a concise overview of the essential elements that make for effective partnering.

Website: www.thepartneringinitiative.org

# 6

# ORGANISATIONAL CHANGE AND SELF-CARE FOR WORKERS



- Organisational change requires a commitment by all levels of management and staff to embrace the implementation of new approaches to treatment.
- Change management is a process that requires planning, coordination and review.
- Audits of current approaches and practices against the proposed change can assist in identifying where energies should be focused and prioritised.
- Conducting a pilot of an intended change and then evaluating the outcomes of a particular approach can assist in staff engagement.
- Professional development and training staff is essential in harnessing commitment to change in approach or practice.
- Promoting a culture of inclusion and support for staff is vital in maintaining best practice approaches with clients. Workplace practices that promote effective supervision and self-care are key parts of a supportive organisational culture.

All the frameworks and practices outlined in this resource have been incorporated into AOD treatment services in Australia and overseas, in varying degrees. Implementing best practices approaches with women accessing AOD treatment requires a whole-of-organisation commitment and implementation process in order for there to be consistency. From policy and procedures through to support structures for workers, there should be evidence of agreement throughout the organisation and its practices.

"It's important to remember that, although training and education are crucial to increasing the capacity of drug and alcohol services in responding to the emerging needs of the community, they are not sufficient on their own to bring about sustained improvement in service delivery and system responses (Roche & Skinner 2005). To bring about sustained improvement through organisational change management, an effective process is essential" (NADA 2013).

The specific steps that make up an effective organisational change process are explored as part of a set of resources developed by NCETA within the *Theory Into Practice Strategies (TIPS) Tool Kits: Organisational Change Management* resource, and are presented here:

#### Stage 1: Unfreezing

Investing time at the start of a change program to prepare and support workers is an essential step to minimise reluctance to change and ensure successful implementation of new work practices.

Issues to be addressed at the unfreezing stage include:

- acknowledging current work practices
- supporting workers' readiness for change
- providing sufficient organisational resources for change
- providing professional development (e.g. education, training) for new work practices
- managing uncertainty associated with change.

#### Stage 2: Changing

Strategies to assist the transition from old to new work practices include:

- conducting trials of change
- engaging in ongoing monitoring and evaluation
- supporting workers to change their behaviour (e.g., support, feedback, rewards, professional development).

#### Stage 3: Confirming and Supporting

Strategies to ensure new behaviours become standard work practice include:

- continuing to offer workplace support for the new work practices
- continuing with monitoring and evaluation of change including making required modifications to the new work practices.

All staff should be involved in assessing and evaluating the current strengths and challenges of the organisation before the change implementation process commences, ensuring all staff have input and take ownership of the change process. A change management plan should incorporate space for staff to discuss challenges, frustrations and success stories, as implementing change can have a significant impact on organisational culture. Clients should be included in the organisational assessment process. This may involve speaking to current and past clients as well as their family and carers.



Find out more...about organisational change management.

Organisational Change. Workforce Development TIPS (Theory Into Practice Strategies): A Resource Kit for the Alcohol and Other Drugs Field (Skinner NCETA 2005).

The Workforce Development (WFD) 'TIPS' Kit provides a unique range of user-friendly and practical WFD tools and resources for the AOD workforce targeted at supervisors, managers, policy makers and other key individuals within the AOD field who are likely to engage in WFD activities and interventions.

# 6.1 Areas for review and enhancement in mixed-gender organisations

Although there is evidence to suggest that women experiencing issues with substance use are best placed in women-only services, there are a number of ways to enhance mixed-gender organisations to better meet the needs of female clients. As indicated in the previous section, organisational change requires commitment and consultation. In order to demonstrate here how a mixed-gender organisation might adopt a family-inclusive approach as an example, the principles of the organisational change model developed by NCETA (2005) will be explored.


# STEP 1: UNFREEZING

One of the first steps in the change management process is to acknowledge the current work practices that fit with family-inclusive approach and/or that work to support the same goals as family inclusive practice within the mixed-gender organisation. It is important to address the potential discomfort workers may feel as part of the change process – just as their clients do in treatment

It is also important at this stage in the process to inform staff of the nature of the changes. In the case of mixed-gender organisations it might be providing the evidence base for family-inclusive practice and the range of supports that will be put in place to support staff in this area – through training, for example. It may also be helpful to have an expert in the area attend the service as motivation. They will be able to respond to staff concerns about how they might incorporate the approach into their work. Providing information in the form of handouts and research literature, diagrams and support tools will also support any training that takes place.

It may also be possible to arrange a site visit to an AOD treatment service where family-inclusive practice has been implemented successfully. This will enable staff to see where the organisation is heading and witness the approach in practice. It may also be helpful to involve staff in considering what is possible in terms of change towards a family-inclusive practice approach.

The framework presented below is a useful starting point in considering what level of organisational change is possible/practicable and the specific areas that require review.



Harnessing the curiosity of staff in applying something that will enhance their practice is the best approach to commencing organisational change and this can be achieved by effective communication, inclusion and support.

# STAGE 2: CHANGING

Implementing actual change is best done in stages and a smaller pilot of the practices might be the best place to begin. For example, after a review of the assessment form and practises, a mixed-gender organisation might decide to trial the new process within a specific timeframe and then evaluate it. This staged approach assists with the change process and allows staff to reflect on how the changes shape their work practices. It may also identify areas for further training and/or support. "Changes that can be tested and evaluated in a trial bases are more likely to be accepted by workers" (Tornatzky & Klein 1982 in Armenikas et al 1999).

A checklist developed by NCETA, Checklist for Family Sensitive Practice in the AOD Field, covers the following elements of organisational practice that might require review:

- 1. Assessment
- 2. Intervention
- **3.** A partnership and empowerment approach that explores the role of clients, families and communities involvement in program development
- 4. Multi-agency and cross-sectoral working
- 5. Workforce development
- 6. Organisational and systems development
- 7. Building leadership and integrated government policy
- 8. Accountability and monitoring

Each element poses some important questions for consideration by the organisation and will help it assess the fit between current practices and what might need altering in order to reflect a family inclusive approach. In practical terms it might be best to formulate a clear plan of staged introductions of new interventions starting with some of the organisational culture aspects first, such as language, policies and procedures. From here, a process of regular audits and evaluations of screening, assessment and treatment planning tools can also be implemented.

An example of some of the questions you might pose when exploring implementing family-inclusive practice is as follows:

- 1. Undertake an organisational checklist or audit in regard to child-friendly practices to ensure that each treatment agency has child-friendly policies and procedures in place. An example of a checklist for this purpose would be one developed by NCETA: Checklist for Family sensitive practice in AOD services.
- 2. Expand education and training of staff to undertake child and parent-sensitive practice.
- **3.** Engage in appropriate clinical supervision for all staff and services where clients have children.
- **4.** Include questions regarding clients' parenting roles and responsibilities as part of a routine assessment and clinical review.

Another useful resource that explores the implementation of family-inclusive practice into AOD treatment, "Responding to the needs of children and parents in families experiencing alcohol and other drug problems" by Gruenert and Tsantefski (2012), provides some key informants' top tips for workers, organisations, and governments and other funders.

# Key informants' top tips for workers

- You do not have to be an expert in family therapy to ask clients about their children or other family members. Be yourself, honest, genuine, open and interested.
- Do not make assumptions. Aim to have some communication with relevant family members and encourage greater communication between family members.
- You probably already have the skills to contact other family members and identify their needs.
- Be aware that trauma and grief may go beyond individuals, and impact families and whole communities.
- Be open and respectful to all you speak with and anticipate some level of anxiety, conflict
  and shame. The stigma of having an AOD problem while being a parent can be enormous.
- Where possible, family members should be informed about their rights and responsibilities, what information will be shared, and how this will be done.
- Provide families with some information and support to address the effects of trauma and build hope for recovery.
- Take a supportive and strengths based approach rather than a punitive or risk focused approach. This will help engagement with parents and improve the likelihood of real change, not simply compliance.
- Include child and parent focused goals in your treatment plan. Think about what children need to be healthy and happy, in addition to any safety issues. Learn more about child development levels and timeframes, and discuss these in supervision and case reviews.
- Setting and monitoring goals is important. So is having a plan B for the care of children.
   As lapses are common in the recovery process, a parent's functioning may vary considerably over time.
- Ensure pregnant women get specialist antenatal support and care.
- Consider inviting a trusted family member or significant person to attend a session with your client. This may be especially important for building trust when meeting families for the first time, and with Indigenous and CALD communities.
- Help parents to understand or explore the possible impact of their AOD use or treatment on their children. Support parents to access material needs and develop new parenting strategies, especially around limit setting for their children.
- Help parents to strengthen their social network.
- Get to know your local child and family support and child protection workers. Know where the maternal and child health and education services are located. Identify child care options.
- Seek and respect children's opinions, but not at the expense of their safety.
- Help children to understand what is happening to their parents. They may need help to understand that family problems are not their fault, and that their parents love them, even if they are unable to care (or adequately care) for them at present.
- Provide children with opportunities to share their experiences with other children in similar circumstances. Tutoring and recreational activities may help children feel normal.
- Older children and young people may need their own intensive and specific long-term support and follow-up.

(Gruenert and Tsantefski 2012)

Within this process of change it is important to set up feedback mechanisms and meet regularly in different forums (all staff, groups of staff, administrative/executive groups) in order to assess how the change process is progressing. This provides an opportunity for staff to be acknowledged and appreciated for the work they are doing. Rewards such as flexi-time, support for professional development or attendance at conferences are also useful in supporting the change process.

# STAGE 3: CONFIRMING/ SUPPORTING

In this final stage, the new approach, family-inclusive practice, is incorporated into standard work practice. "The 'new way' becomes the 'normal' way" (NCETA 2005:15).

It is important in this stage to continue to provide support to workers and to maintain regular review, evaluation and feedback.



Find out more...about Organisational Change.

Organisational Change. Workforce Development TIPS (Theory Into Practice Strategies): A Resource Kit for the Alcohol and Other Drugs Field. NCETA 2005 (Skinner, N In N. Skinner, A.M. Roche, J. O'Connor, Y. Pollard, & C. Todd (Eds.).

Website: www.nceta.flinders.edu.au/workforce

The Resource Kit comprises 14 chapters: an introduction to workforce development and 13 workforce development topics relevant to the AOD field. Each chapter contains evidence-based strategies to address a particular workforce development issue, as well as resources and tools that can be used to implement the strategies. Each chapter can be treated as a stand alone section, however, as workforce development topics are inherently interrelated, links between chapters are identified throughout the Kit. Organisational Change is the 7th chapter in the Resource Kit.

# 6.2 Enhancing organisational approaches that promote self-care

Working in the AOD treatment sector can be very rewarding. However, it is an accepted part of the work that clients who present for treatment are complex and that problematic substance use is a chronic, relapsing condition. What this means in practice is that without comprehensive support and supervision, the challenges of working with AOD clients can become overwhelming – leading to stress and potential burnout.

As we've seen, women who present for AOD treatment can frequently experience domestic violence and trauma, legal concerns, child protection issues and difficulty with finding adequate housing. These challenges, when coupled with negative attitudes within the community towards women who use drugs, can lead to an increased need for workers to advocate for their clients. As such, workers must sometimes bear the brunt of negative attitudes expressed towards them personally. In order to establish a culture of self-care in an AOD treatment organisation there needs to be a range of options that engage AOD workers to reflect on their practice, learn new skills and debrief about their concerns.

A number of strategies can be employed by an AOD treatment organisation to enhance worker self-care. They can include:

- peer supervision
- external clinical supervision
- training and professional development
- team building exercises and leisure activities
- access to a robust Employee Assistance Program.

NCETA suggests a two-pronged approach to addressing stress and burnout in drug and alcohol services (Skinner 2005):

- 1. There should be organisational strategies that focus on altering the work environment and/or the conditions causing the stress and burnout.
- **2.** The individual worker should focus on developing coping strategies and stress management techniques.

The following table highlights the strategies services can implement to provide workplace support.

	SOCIAL/EMOTIONAL SUPPORT	INSTRUMENTAL SUPPORT
Organisation	<ul><li>Ensuring fairness of treatment</li><li>Providing valued rewards</li><li>Ensuring supportive supervision</li></ul>	<ul> <li>Ensuring good job conditions (physical safety, job security, promotion paths, autonomy)</li> <li>Addressing work overload</li> <li>Addressing role ambiguity or conflict</li> <li>Providing access to high-quality resources and equipment</li> </ul>
Managers/ supervisors	<ul> <li>Channelling/facilitating organisational support</li> <li>Providing positive social interaction (praise, encouragement, caring, respect)</li> <li>Recognising and rewarding good work</li> <li>Involving workers in decision-making</li> </ul>	
Co-workers	<ul> <li>Providing positive social interaction (praise, encouragement, caring, respect)</li> </ul>	<ul> <li>Providing help and advice</li> <li>Filling in when others are absent</li> <li>Assisting with heavy workloads</li> <li>Providing constructive feedback</li> <li>Appreciation and recognition</li> <li>Sharing duties and responsibilities</li> </ul>



Find out more...about AOD worker self-care and support.

**NCETA Workforce Development 'TIPS'** (Theory Into Practice Strategies): A Resource Kit for the Alcohol and Other Drugs Field (NCETA 2005).

This toolkit has a two specific sections that focus on Worker Wellbeing and Worker Support. Website: www.nceta.flinders.edu.au/workforce

# The Feeling Deadly-Working Deadly Indigenous Wellbeing Toolkit NCETA 2013

**The** 'Feeling Deadly, Working Deadly' Resource Kit is aimed at reducing stress and burnout, and enhancing wellbeing amongst Aboriginal and Torres Strait Islander alcohol and other drug (AOD) workers. It forms part of NCETA's work on Aboriginal and Torres Strait Islander Worker Wellbeing. It was funded by the Australian Government Department of Health.

**Counselling guidelines: Alcohol and other drug issues** 3rd Edition WA Drug and Alcohol Authority (Marsh et al. 2013)

These guidelines includes over 50 chapters exploring some of the key skills needed to work at an individual level with people who have AOD problems including a case management, group work, referral, working with intoxicated and incarcerated clients, clinical supervision and stress and burnout.

# 7

# RESOURCES AND GUIDELINES

The resources and guidelines section provides a brief selection of information and support services, resources and toolkits for workers, and government produced guidelines and strategies which relate to working with women engaged in the AOD sector. It is by no means an exhaustive list, however many of the websites noted here provide further useful links to an expansive array of support documents and resources.

# 7.1 Information and support

# ALCOHOL AND OTHER DRUG SUPPORT

# Alcohol and Drug Information Service (ADIS)

ADIS is a 24-hour helpline which provides alcohol and drug related information, support, crisis counselling and referral to NSW services.

Phone: 02 9361 800 or 1800 422 599 Website: www.yourroom.com.au

# Alcohol, Tobacco and other Drugs Association of ACT (ATODA)

ATODA is the peak body representing the non-government and government alcohol, tobacco and other drug (ATOD) sector in the ACT.

Website: www.atoda.org.au

# Alcohol, Tobacco and other Drugs Council of Tasmania (ATDC)

The ATDC is the peak body representing the interests of the community service organisations that provide services to people with substance misuse issues in Tasmania.

Website: www.atdc.org.au

# Association of Alcohol and other Drugs Agencies in Northern Territory (AADANT)

AADANT is the peak body for the non government alcohol and other drug sector in the Northern Territory.

Website: www.aadant.org.au

# **Drugfields**

Drugfields aims to support the Australian alcohol and drug workforce through a professional development service and website which provide the assistance to individuals and organisations.

Website: www.drugfields.org.au

# Drug and Alcohol Multicultural Education Centre (DAMEC)

DAMEC was established in 1989 as an access and equity initiative designed to promote access by culturally and linguistically diverse (CALD) clients to alcohol and other drug services. DAMEC provide counselling and support for CALD people and have a research and information focus. DAMEC works within a harm minimisation framework, and uses the term alcohol and other drugs to describe both licit and illicit drugs, including alcohol, tobacco and prescription medicines that are used outside medical advice.

Phone: 02 9699 3552

Website: www.damec.org.au

# Drug and Alcohol Specialist Advisory Service (DASAS)

DASAS is a free alcohol and drug-related advisory telephone service for doctors, nurses, and other health professionals. Specialist medical consultants are on-call 24 hours to provide advice on diagnosis and management of patients. DASAS is especially designed to support regional and rural areas in NSW, but is available to any health professional.

Phone: 02 9361 8006 or 1800 023 687

# **Family Drug Support**

Family Drug Support (FDS) was formed in 1997 after its founder, Tony Trimingham lost his son to a heroin overdose. FDS is a caring, non-religious and non-judgemental organisation, primarily made up of volunteers who have experienced firsthand the trauma and chaos of having family members with drug dependency. FDS provides a 24 hour/7 day a week telephone support line, website, support groups and courses for family members and significant others.

Phone: 1300 368 136 Website: www.fds.org.au

### Foundation of Alcohol and Research Education (FARE)

FARE is an independent, not-for-profit organisation working to stop the harm caused by alcohol.

Website: www.fare.org.au

# Network of Alcohol and other Drugs Agencies (NADA)

NADA is the peak body for the non government alcohol and other drugs sector in NSW.

Website: www.nada.org.au

### **NSW Users and AIDS Association (NUAA)**

NUAA is the peak drug user organisation in NSW. It is a not-for-profit organisation advocating for people who use drugs, particularly those who inject drugs. NUAA provides education, practical support, information and advocacy to users of illicit drugs and their friends and allies.

Website: www.nuaa.org.au

# Queensland Network of Alcohol and Drug Agencies (QNADA)

QNADA is the peak organisation representing the views of the non-government alcohol and other drug sector in Queensland.

Website: www.qnada.org.au

# South Australian Network of Drug and Alcohol Agencies (SANDAS)

SANDAS is the peak body providing independent, statewide representation, advocacy and support for non-government organisations working in the South Australian alcohol and other drug sector, through networking and policy development.

Website: www.sandas.org.au

### Victorian Alcohol and Drug Association (VAADA)

VAADA is the peak body representing drug and alcohol services in Victoria; it provides leadership, representation, advocacy and information to both drug-and-alcohol and non drug and alcohol related sectors.

Website: www.vaada.org.au

# Western Australian Network of Alcohol and other Drug Agencies (WANADA)

WANADA is the peak body for the drug and alcohol education, prevention, treatment and support sector in Western Australia.

Website: www.wanada.org.au

# COMPLEX NEEDS

# **COMPLEX** Acquired Brain Injury Services

A not-for-profit specialist service for people with an aquired brain injury (ABI). Its aim is to enhance people's lifestyle by maintaining and promoting independence and providing opportunities for socialisation and community integration with appropriate and quality service provision.

Website: www.abis.org.au

# Ageing Disability and Homecare (ADHC) ABI Web Resource

A webpage dedicated to ABI on the Family and Community Services Ageing Disability & Home Care website provides information on care and support pathways for people with an ABI, as well as information on the new ADHC training resource (web-based and face-to-face training options available).

Website: www.abistafftraining.info

### arbias

arbias provides support to people with an acquired brain injury, specialising in alcohol and other drug related brain injury in NSW and Victoria. arbias is a case management, specialist assessment and intervention service targeted at people living with an alcohol-related acquired brain injury aged 16 to 65. arbias also provides a comprehensive range of training package options. Its training program provides practical strategies for working with people with acquired brain impairment and associated complex support needs.

Website: www.arbias.org.au

# Brain Injury Australia (BIA)

BIA is the peak ABI advocacy body representing, through its state and territory member organisations, the needs of people with an ABI, their families and carers. It works at a national level to make sure all people living with ABI have access to the supports and resources they need to optimise their social and economic participation in the community.

Website: www.braininjuryaustralia.org.au

### Community Restorative Centre (CRC) NSW

CRC is a NSW community organisation dedicated to supporting people affected by the criminal justice system, particularly prisoners, ex-prisoners, and their families and friends. Staff offer personal and practical assistance such as counselling, accommodation, a subsidised transport service, a court support service, outreach to prisons, information, advice and referrals.

Website: www.crcnsw.org.au

### Fetal Alcohol Spectrum Disorder and Justice Website

This Canadian site is designed for justice system professionals and others who want to understand more about FASD. It provides information and resources about FASD, including background information, case law, legal resources and strategies for effective intervention.

Website: www.fasdjustice.on.ca

# National Organisation for Fetal Alcohol Spectrum Disorder (NOFASD)

NOFASD Australia is an independent not-for-profit charitable organisation. NOFASD are the national peak organisation representing the interests of individuals and families living with Fetal Alcohol Spectrum Disorders (FASD). The website provides information, training and support related to FASD.

Website: www.nofasd.org.au

# FAMILY AND DOMESTIC VIOLENCE

### The Domestic Violence Line

This service provides telephone counselling, information and referrals for women and same-sex partners who are experiencing or have experienced domestic violence. Trained female caseworkers are sensitive to the needs of people who have experienced domestic violence. Domestic Violence Line staff are aware of the special needs of Aboriginal women and women from other cultures, as well as those living in rural and remote areas. Interpreters and TTY can be arranged where necessary to ensure that all people, regardless of their language or disability can use the service. The service has an extensive list of contacts, people and services across NSW who can help. For information, support and referrals regarding FDV contact the Domestic Violence 24 Hour Hotline.

Phone: 1800 656 463

**Domestic Violence NSW** (DVNSW) is the peak, state-wide representative body for a range of specialist domestic and family violence services.

Website: www.dvnsw.org.au

**Domestic Violence Service Management** (DVSM) aim to prevent domestic and family violence and homelessness, and support recovery. They provide professional services to the community services sector and client service delivery based on a social justice framework.

Website: www.dvnswsm.org.au

# **Domestic Violence Website**

Includes information regarding FDV including telephone referral lines, tips and apps that can be downloaded, providing information for those experiencing FDV and/or those experiencing it.

Website: www.domesticviolence.nsw.gov.au

# MENTAL HEALTH

# **Beyond Blue**

Beyond Blue is an independent, not-for-profit organisation working to increase awareness and understanding of anxiety and depression in Australia and to reduce the associated stigma. They offer support via a 24 hour/7 day a week helpline and also offer web chat services.

Phone: 1300 22 4636

Website: beyondblue.com.au

# **Dual Diagnosis Australia and New Zealand**

Dual Diagnosis Australia and New Zealand is a resource repository created to contribute to better outcomes for persons with co-existing substance use and mental health disorders.

Website: www.dualdiagnosis.org.au/home

### Mental Health Association

Mental Health Association runs both the Mental Health Information Service and the Anxiety Disorders Information Line. These lines provide information, telephone support and referral for issues relating to mental health.

Mental Health Information Service - Phone 1300 794 991

Website: www.mentalhealth.asn.au

# Mental Health Coordinating Council (MHCC)

The MHCC is the peak body for community mental health organisations in NSW. MHCC provides leadership and representation to its membership and seeks to improve, promote and develop quality mental health services for the community. MHCC provides a range of training and education through its Learning and Development Unit.

Website: www.mhcc.org.au

# WOMEN'S HEALTH

# Aboriginal women and sexual assault: 'Hey sis, we've got your back'

The Aboriginal Women's Sexual Assault Network was borne out of collaboration between Mudgin-Gal Aboriginal Corporation and Rape and Domestic Violence Services Australia (formerly NSW Rape Crisis Centre). Rape and Domestic Violence Services Australia is currently managing the initiative, with the assistance of an Advisory Committee of respected Aboriginal women.

The network seeks to bring together Aboriginal women who are working against sexual assault in their communities, or who would like to stand strong and support others in their communities who have experienced or been impacted by sexual assault. Through sharing stories and knowledge of what works and what doesn't, through the provision of culturally appropriate training, and through working together to develop and implement initiatives to prevent sexual assault at the community level, the network aims to reduce the rates of sexual assault in Aboriginal communities.

Website: www.heysis.com.au

### **ACON**

ACON provide specific support services that relate to the health and wellbeing of Lesbian, Bisexual and Transgender women. In addition, through partnerships with academic institutions, ACON have engaged with significant research projects aimed at improving health outcomes for the LGBTI community as a whole.

Website: www.acon.org.au

### Australian Women's Health Network

The Australian Women's Health Network is a health promotion advocacy organisation that provides a national voice on women's health, based on informed consultation with members. Through the application of a social view of health, it provides a woman-centred analysis of all models of health and medical care and research. It maintains that women's health is a key social and political issue and must be allocated adequate resources to make a real difference.

It aims to foster the development not only of women's health services but of stronger community-based primary health care services generally, which it sees as essential to improve population health outcomes. It advocates collaboration and partnership between relevant agencies on all issues affecting health.

Website: www.awhn.org.au

### Women's Health NSW

Women's Health NSW (WHNSW) is an association of statewide women's health centres and specialist women's centres. All centres are non-government, community based, feminist services that provide choices for women to determine their individual health needs. WHNSW centres aim to improve the health status of women by providing a unique, holistic, womancentred approach to primary health care. The centres aim to blend medical and clinical services and a range of counselling, health promotion, education, self-help and consumer advocacy services. They also aim to provide women with the knowledge, skills and resources to enable women to take more responsibility over factors the adversely affect their health. Website: www.whnsw.asn.au

# 7.2 Practice resources and toolkits

# RESOURCES

# ABORIGINAL Aboriginal-specific Community Reinforcement Approach (CRA) Training Manual NDARC (Rose et al. 2014).

The Community Reinforcement Approach (CRA) as it is described in this manual has been adapted from a comprehensive explanation as published in Meyers and Smith (1995) Clinical Guide to Alcohol Treatment: The Community Reinforcement Approach. This adapted CRA manual is designed for Indigenous Health and Family Workers and other Workers within Australian Indigenous community settings to support people who are at risk for alcohol related harm.

# Australian Indigenous Health Infonet

A comprehensive web resource for working effectively and supportively with Indigenous Australians that provides information, training resources, projects, latest news and links. A web resource that is a 'one-stop info-shop' for people interested in improving the health of Indigenous Australians.

Website: www.healthinfonet.ecu.edu.au

# Implementing a Soft Entry Approach to AOD intervention with Aboriginal women Lyndon Community 2014

This is a PowerPoint presentation outlining a unique and effective approach used by the Lyndon Community to make contact with Aboriginal women with the aim of potentially providing alcohol and other drug support.

# National Aboriginal and Torres Strait Islander Women's Alliance

The National Aboriginal and Torres Strait Islander Women's Alliance (NATSIWA) was established in 2009 to empower Aboriginal and Torres Strait Islander Women to have a strong and effective voice in the domestic and international policy advocacy process.

Website: www.natsiwa.org.au

# NIDAC AOD Treatment Report NIDAC 2014

This paper was developed by the National Indigenous Drug and Alcohol Committee (NIDAC) in response to a misperception that effective AOD treatment is not available for Aboriginal and Torres Strait Islander peoples. The paper aims to allay these misperceptions by outlining who can benefit from receiving treatment; what treatment is known to work; key principles that should guide the application of treatment; and what constitutes effective treatment for Aboriginal and Torres Strait Islander people.

Website: www.nidac.org.au

Strengthening Aboriginal Family Functioning: What works and why? Child Family Community Australia, Australian Institute if Family Studies 2014 (Lohoar, Bureta & Kennedy)

This paper explores and discusses the contemporary evidence base to provide insights into the protective effects and risks that influence forms of functioning among Aboriginal families. It also has a section on specific resources.

### NADA RESOURCES

# Complex Needs Capable: A Practice Resource for Drug and Alcohol Services NADA 2013

This resource provides practice tips, tools and templates that can support alcohol and other drugs staff and organisations working with clients with complex needs such as acquired brain injury, intellectual disability, fetal alcohol spectrum disorders and contact with the criminal justice system.

Website: www.nada.org.au /www.complexneedscapable.org.au

# From Individuals to Families: Single Session Consultations (DVD) NADA 2012

This DVD, developed in partnership with NADA and The Bouverie Centre, provides a practical case example from an alcohol and other drug service. It also included interviews with staff from three NADA members discussing the implementation of the single session approach to working with families in alcohol and other drug settings.

# Tools for Change: A New Way of Working with Families and Carers NADA 2009

Tools for change provides a range of interventions, practice tips, service models, resources and training organisations to assist services in working with families.

Website: www.nada.org.au

# Working with Diversity in Alcohol and other Drug Settings NADA 2014

The resource has been developed to support non government alcohol and other drug services work with the diversity of clients that access our services and represent the NSW population. It contains examples of best practice.

A supplementary document: **Working with Diversity in Alcohol and other Drug Settings - Useful Resources** NADA 2015 also provides a comprehensive list of updated resources of use.

NADA recognises that there are other populations that have not been included such as young people, women, children, and those with a disability. NADA hope to update the guide as information is developed. For best practice examples in working with young people, access the Dovetail Youth Alcohol and Drug Good Practice Guide.

Website: www.nada.org.au

# FAMILY AND DOMESTIC VIOLENCE

# Can I ask...? An alcohol and drug clinician's guide to addressing family and domestic violence NCETA 2013 (White et al.)

Can I ask...? explores the relationship between AOD and FDV, with a focus on identifying how the AOD sector can better support clients who have co-existing AOD and FDV issues, and minimise associated harms experienced by their children.

The **Domestic Violence Resource Centre Victoria** (DVRCV) provides training, publications, research and resources to those experiencing (or who have experienced) family violence, and practitioners and service organisations who work with family violence survivors. Website: www.dvrcv.org.au

Supporting Women with Complex Needs: The relationship between substance use and domestic and family violence Women's Council for Domestic and Family Violence Services (WA) and WANADA 2009

This resource explores how service providers can work in collaborative and respectful ways with women who have experienced violence and who have substance use concerns. It identifies how common it is for women to experience both concerns, and how critical is it for service providers to assist women to make the links and to determine paced and achievable paths to change, growth and safety.

# Walking a Tightrope: Alcohol and other drug use and violence: A guide for families NCETA 2014

Walking a Tightrope is a comprehensive resource for people who have a family member who uses both alcohol and other drugs (AOD) and violence in their relationships. The resource has been produced through a partnership between Family Drug Support (FDS) and the National Centre for Education and Training on Addiction (NCETA) and was funded by the Department of Social Security.

# FAMILY INCLUSIVE PRACTICE

### Children of Parents with a Mental Illness Website

The COPMI (Children of Parents with a Mental Illness) national initiative develops information for parents, their partners, carers, family and friends in support of these children. This information includes online training courses developed by COPMI for professionals to support families either individually or through community services and programs.

Website: www.copmi.net.au

# Familiar Needs; Working with Children and Families VAADA 2011

Resource developed by VAADA to provide AOD treatment services with a snapshot of resources available and practice tips for working with families and children.

# Families and friends affected by the drug and alcohol use of someone close NSW Health 2004

This resource answers common questions and concerns of those who are experiencing the challenges of having someone they care about engage in problematic substance use.

# Family Focus Project Toolkit Eastern Drug and Alcohol Service 2010

The Family Focus Toolkit is a collection of selected resources including screening tools, questionnaires, worksheets, and utility practice tools gathered from the sector, research and professional bodies.

# For Kids' Sake: A Workforce Development Resource for Family Sensitive Policy and Practice in the Alcohol and Other Drugs Sector NCETA 2010 (Battams et.al.)

For Kids' Sake, A Family Sensitive Policy and Practice Toolkit aims to minimise cases of child abuse or neglect by increasing collaboration between child and adult service agencies. It is a new initiative developed by NCETA to improve the safety and welfare of children with parents who misuse alcohol or drugs. The toolkit builds a bridge between the alcohol and other drugs treatment and child protection sectors to improve cooperation and collaboration.

# From individuals to families: A client centered framework for involving families, The Bouverie Centre, La Trobe University

This document, was developed for adult mental health and alcohol and drug services providers as a framework for a comprehensive service response to a client's family, social network or kinship group in the context of client-centred care and sets out key issues in implementing this framework. The ideas presented can guide both individual services and groups of services within a catchment area in determining how they can best involve and respond to the needs of families.

# From Individuals to Families: Single Session Consultations (DVD) NADA and the Bouverie Centre 2012

This DVD, developed in partnership with NADA and The Bouverie Centre, provides a practical case example from an alcohol and other drug service. It also included interviews with staff from three NADA members discussing the implementation of the single session approach to working with families in alcohol and other drug settings.

# Legal Aid Care and Protection Factsheets

These fact sheets provide information that helps people navigate the Children's Court and children's services. The five factsheets in the series cover the process of going to the Children's Court and the role of Family and Community Services.

The *Kids in Care* fact sheets have been translated into 10 community languages – Arabic, Bengali, Dari/Farsi, Dinka, Kirundi, Chinese (Sim), Swahili, Tamil, Thai and Vietnamese. They can be **ordered** and are also available online.

Website: www.legalaid.nsw.gov.au/publications/

# **The signs of safety: Child protection practice framework** Department of Child Protection Western Australia 2011

Is a policy document that seeks to create a more constructive culture around child protection organisation and practice. Central to this is the use of specific practice tools and processes where professionals and family members can engage with each other in partnership to address situations of child abuse and maltreatment.

# Tools for Change: A New Way of Working with Families and Carers NADA 2009

Tools for change provides a range of interventions, practice tips, service models, resources and training organisations to assist services in working with families.

Website: www.nada.org.au

# CALD COMMUNITIES

# Drug and Alcohol Multicultural Education Centre (DAMEC)

DAMEC was established in 1989 as an access and equity initiative designed to promote access by culturally & linguistically diverse (CALD) clients to alcohol and other drug services. DAMEC provide counselling and support for CALD people and have a research and information focus. DAMEC works within a harm minimisation framework, and uses the term alcohol and other drugs to describe both licit and illicit drugs, including alcohol, tobacco and prescription medicines that are used outside medical advice.

# Helping asylum seeker and refugee background communities with problematic alcohol and other drug use: A guide for community support and AOD workers QNADA 2015 (Pont B and Jaworski A)

The guide is divided into two sections, the first is for workers supporting people from refugee backgrounds across the general health and community services sector who may not have specific training or experience in alcohol and other drugs service provision. And the second is for those working in the AOD sector and discusses issues that are relevant across the range of treatment settings.

# **Immigrant Women's Speakout Association NSW**

Immigrant Women's Speakout Association is the peak advocacy, information/referral and research body representing the ideas and issues of immigrant and refugee women in NSW. The Association also undertakes community development projects and provides direct services including in the areas of domestic violence and employment, education and training. Speakout is a community-based organisation, managed by women of non-English speaking background. There are a number of resources available via the website, including training opportunities.

Website: www.speakout.org.au

# Respect: Best practice approaches for working with culturally diverse clients in AOD treatment settings. DAMEC 2014

Website: www.damec.org.au

### Perinatal Mental Health for Women from CALD Backgrounds (Beyond Blue)

This is a brief guide for health care workers that covers a number of issues and barriers CALD women experience in accessing support during pregnancy, birth and in relation to parenting. It attends to the needs of CALD women and how health services can be more responsive in relation to the experience of pregnancy, birth and parenting.

Website: www.beyondblue.org.au

# **Talking therapies for people with problematic substance use** Te Pou o Te Whakaaro Nui 2010

This guide provides useful background information on problematic substance use, suggestions for interventions during the various stages of treatment, and information on the range of talking therapies currently utilised by practitioners in New Zealand. Also includes sections on Maori, other Pacific Islander cultures and working with refugees.

# LGBTI COMMUNITIES

# LGBTI Best Practices for Lesbian/Bisexual Women with Substance Use Disorders (Eliason 2009)

This paper reviews what we know about substance abuse patterns among lesbian and bisexual women and suggests best practices.

### **GLBTIQ** Domestic Violence Toolkit

A resource for agencies and service providers supporting GLBTI victims of domestic violence and their families.

Website: www.avp.acon.org.au/anti-violence/resources/GLBTIQDV-toolkit

### Lesbian Relationships and Abuse - Domestic Violence Resource Centre Victoria

Help and advice on the DVRCV website that provides information regarding the experience of domestic violence within lesbian relationships.

Website: www.dvrcv.org.au

# TRANSGENDER WOMEN

### The Gender Centre

The Gender Centre offers a wide range of services to transgender and gender diverse people, their partners, families and friends in NSW. They also act as an education, support, training and referral/resource centre to other organisations and service providers. The Gender Centre aim to provide a high quality service which acknowledges human rights and ensures respect and confidentiality.

Website: www.gendercentre.org.au

**The First Australian National Trans Mental Health Study: Summary of Results** Curtin University 2013 (Hyde Z, et al)

This report provides an overview of the findings of The First Australian National Trans Mental Health Study. The study was designed to investigate the mental health and wellbeing of trans people living in Australia.

### COMPLEX NEEDS

# Complex Needs Capable: A Practice Resource for Drug and Alcohol Services NADA 2013

This resource provides practice tips, tools and templates that can support alcohol and other drugs staff and organisations working with clients with complex needs such as acquired brain injury, intellectual disability, fetal alcohol spectrum disorders and contact with the criminal justice system.

Website: www.nada.org.au /www.complexneedscapable.org.au

**Counselling guidelines: Alcohol and other drug issues 3rd Edition** WA Drug and Alcohol Authority (Marsh et al. 2013)

These guidelines includes over 50 chapters exploring some of the key skills needed to work at an individual level with people who have AOD problems including a case management, group work, referral, working with intoxicated and incarcerated clients, clinical supervision and stress and burnout.

Guidelines on the Management of Co-Occurring Alcohol and Other Drug and Mental Health Conditions in Alcohol and Other Drug Treatment Settings NSW Department of Health 2009

Available from: www.ndarc.med.unsw.edu.au/

**Mental Health Reference resource for Drug and Alcohol Workers** NSW Department of Health 2007

Available from NSW Ministry of Health: www.health.nsw.gov.au

# MHCC Trauma Informed Care and Practice webpage

This website contains a range of resources and events listings and a network on traumainformed care and practice.

Website: www.mhcc.org.au

NSW Clinical Guidelines for the Care of Persons with Comorbid Mental Illness and Substance Use Disorders in Acute Care Settings NSW Department of Health 2009

Available from NSW Ministry of Health: www.health.nsw.gov.au

# Pathways of recovery (Rickwood 2005)

This monograph raises the vital issue of the role of relapse prevention in the recovery process for people who have been seriously affected by mental illness. While preventing further episodes of mental illness should be a routine component of treatment and continuing care for people with mental illness, feedback from consumers and carers shows that frequently this is not the case. Too often the experience of consumers, and their families and carers, is of a crisis-focussed mental health system that doesn't respond early enough to avert further episodes, nor prioritise rehabilitation and relapse prevention as essential components of ongoing continuing care to facilitate recovery. Although relapse prevention has traditionally been viewed as an illness focussed approach, it is reconceptualised here as one of the essential components of a recovery-oriented mental health system.

# PREGNANCY AND PARENTING

# It's time to have the conversation: Understanding the treatment needs of women who are pregnant and alcohol dependent. (NDARC 2013).

This report presents a narrative literature review of treatments available to pregnant women who have alcohol use disorders and findings from interviews with key stakeholders regarding current treatment practices and areas requiring improvement.

# Supporting Pregnant Women who Use Alcohol or other Drugs: A guide for primary health professionals NDARC 2014.

Health care professionals can make a substantial difference to the health of women and their babies by identifying and supporting women who use alcohol or other drugs during pregnancy. This guide is intended for a range of health professionals, in a variety of settings to help support and provide information to pregnant women who use alcohol and other drugs. It also contains a detailed resource list of additional support services and resources.

# What Family Sensitive Practice Means for Alcohol and Other Drug Workers: A Survey Report NCETA 2010 (Trifnoff et al.)

The **report** is the first of a range of publications being developed by NCETA to address the issue of child and parent-sensitive practice. The initial phase of this project involved a survey of the drug and alcohol workforce to identify current work practices, individual knowledge and attitudes and organisational policies and support in relation to child and parent-sensitive work practice. This report presents the results of the survey and outlines.

# TRAUMA INFORMED CARE

### Adults Surviving Child Abuse (ASCA)

ASCA is a national organisation that works to improve the lives of adult survivors of child abuse throughout Australia. ASCA offers a number of training courses and has resources suitable for professionals working with adult survivors of child abuse.

# ASCA's Practice Guidelines for Treatment of Complex Trauma and Trauma Informed Care and Service Delivery 2013

ASCA's Practice Guidelines for Treatment of Complex Trauma and Trauma Informed Care and Service Delivery present the collective wisdom of the last two decades of national and international research in the trauma field. They are poised to revolutionise possibilities for recovery for the large numbers of people with unresolved "complex trauma" - child abuse in all its forms, neglect, family and community violence and other adverse childhood events. They establish insights that optimism about recovery from complex trauma is warranted, and that childhood trauma can be resolved.

# Website: www.asca.org.au

### **Education Centre Against Violence (ECAV)**

ECAV is a state-wide unit responsible for training programs in the specialised areas of adult and child sexual assault, domestic and Aboriginal family violence and physical and emotional abuse and neglect of children across NSW. ECAV provides the mandated training for specialist child protection, sexual assault and Aboriginal family health workers, as well as targeted training for mental health and drug and alcohol workers in government and non-government organisations Website: www.ecav.health.nsw.gov.au

# Guidelines for Trauma-Informed Practices in Women's Substance Use Services Jean Tweed Centre for Women and their families 2013

This Canadian resource sets out to specifically focus on the intersections of trauma and substance use issues among adult women. It provides background information and best practice approaches to working with women engaged in substance use services from a trauma-informed approach to practice.

# NSW Service for the Treatment and Rehabilitation of Torture and Trauma Survivors (STARTTS)

STARTTS supports refugees recover from their experiences and build a new life in Australia. STARTTS is committed to assisting and resourcing people and organisations to provide appropriate and culturally sensitive services to refugee survivors of torture and trauma. Additionally, STARTTS provides a range of counselling and therapeutic services. Website: www.startts.org.au

# Trauma-Informed Organizational Toolkit for homelessness services (Guarino et al. 2009)

This toolkit offers homeless service providers with concrete guidelines for how to modify their practices and policies to ensure that they are responding appropriately to the needs of families who have experienced traumatic stress. AOD services can also use this resource to assist them in becoming trauma informed.

# 7.3 Government guidelines and strategies

**Child Wellbeing and Child Protection – NSW Interagency Guidelines** NSW Department of Family and Community Services 2006

These guidelines were produced to support collaborative efforts to improve child wellbeing and child protection in NSW. They cover inter-agency collaboration initiatives; roles and responsibilities of agencies in child protection matters; making a child protection report voluntarily or through mandatory reporting requirements; exchange of information with FaCS; criminal proceedings; and best practice in working with children and families.

# Clinical Guidelines for the Management of Substance Use During Pregnancy, Birth and the Postnatal Period 2014

These guidelines contain information on the use of buprenorphine maintenance treatment during pregnancy; cannabis use in pregnancy; a recommendation for abstinence from alcohol during pregnancy; and updated information on child protection legislation in NSW. New sections include prescription opioids use in pregnancy; use of antidepressants in pregnancy; management of withdrawal; management of incarcerated pregnant women; and management of women in rural and remote areas.

# Keep Them Safe: A shared approach to child wellbeing

In response to findings from the Wood Special Commission of Inquiry into Child Protection Services in NSW, the NSW government enacted a five year plan entitled Keep Them Safe, which spans the years 2009-2014. Keep Them Safe was designed to enhance the broader service system in NSW to improve prevention and early intervention services; protect children at risk; support Aboriginal children and families; and to strengthen partnerships with non government agencies to improve service delivery to vulnerable families. As a result of Keep Them Safe, a number of guidelines and legislative changes have taken place in the child protection arena in NSW, which are described briefly below.

Website: www.keepthemsafe.nsw.gov.au

# Neonatal Abstinence Syndrome Guidelines NSW Ministry for Health 2013

These guidelines outline the minimum standards for the management of babies born to mothers with a history of opioid use or dependence, including women receiving opioid substitution therapy or using prescription pharmaceutical opioids. The guidelines cover antenatal care, the management of withdrawal syndromes in neonates, and postnatal care.

# **NSW Health Drug and Alcohol Psychosocial Interventions: Professional Practice Guidelines** NSW Dept. of Health 2008

The Drug and Alcohol (D&A) Psychosocial Interventions Professional Practice Guidelines are the first generic professional guidelines for psychosocial interventions to be developed in NSW for drug and alcohol practice. The Guidelines take a stepped care approach to drug and alcohol treatment, which focuses on the adoption of best practice models for people with drug and alcohol issues, and reflects current best evidence and practice in the drug and alcohol counselling field.

# NSW Health Framework for Women's Health 2013 NSW Ministry of Health 2013

The Women's Health Framework has been developed to focus action not only within the Health system, but also promotes opportunities for the co-ordination and delivery of NSW Government services and programs with the potential to enhance the health of all women across the state. This Framework brings together these aspects – highlighting the different needs of populations of women, the evidence for investment in health priorities, and the framework for collaboration across government and non-government sectors to deliver appropriate services and programs for the diverse needs of women in all of our communities.

**NSW Mandatory Reporter Guide** NSW Department of Family and Community Services 2014 All non government drug and alcohol service providers in NSW are mandatory reporters under child protection legislation. To assist workers to navigate the 'grey areas' and make confident decisions when considering making a child protection report to Family and Community Services.

# Supporting Families Early Package – SAFE START Guidelines: Improving mental health outcomes for parents and infants NSW Department of Health 2009

Safe Start is one of a suite of three documents aimed at integrating care for women, infants and families in the perinatal period. This document provides guidance on conducting psychosocial assessment, risk prevention and early intervention. Strategies to coordinate clinical responses to issues identified during assessment are also suggested, including effective responses to parental mental health problems and perinatal psychosocial issues, as well as advice on assisting mothers that have problems with substance use.

# 7.2 Training and workforce development centres

Adults Surviving Child Abuse (ASCA): www.asca.org.au

Centre for Community Welfare Training (CCWT): www.acwa.asn.au/CCWT

Education Centre Against Violence (ECAV): www.ecav.health.nsw.gov.au

Family Worker Training & Development Program Inc: www.fwtdp.org.au

Intellectual Disability Rights Service (IDRS): www.idrs.org.au

Mental Health Coordinating Council (MHCC), Learning & Development Unit: www.mhcc.org.au/learning-and-training/default.aspx

NSW Ombudsman: www.ombo.nsw.gov.au/trainingworkshops/index.html

NSW Institute of Psychiatry (NSWIOP): www.nswiop.nsw.edu.au

Uniting Care Institute of Family Practice: www.ifp.nsw.edu.au

# 8 | REFERENCES

ACON Strategic Plan 2013-2018. NSW Health.

Adams SM, Peden AR, Hall LA, Rayens MK, Staten RR and Leukefeld CG. Predictors of retention of women offenders in a community-based residential substance abuse treatment program. Journal of Addictions Nursing 2011.22(3):103-16.

Albrecht J, Lindsay B and Terplan M. Effect of waiting time on substance abuse treatment completion in pregnant women. Journal Substance Abuse Treatment 2011.41(1):71-7.

Allan J and Campbell M. Improving access to hard-to-reach services: a soft entry approach to drug and alcohol services for rural Australian Aboriginal communities. Social Work Health Care 2011.50(6):443-65.

Allan J and Kemp M. Aboriginal and non aboriginal women in New South Wales non government organisation (NGO) drug and alcohol treatment and the implications for social work: Who starts, who finishes, and where do they come from? Australian Social Work 2011.64(1):68-83.

Al-Otaiba Z, Epstein EE, McCrady B and Cook S. Age-based differences in treatment outcome among alcohol-dependent women. Psychology Addictive Behaviors 2012.26(3):423-31.

Amato L MS, Davoli M, Vecchi S, Ferri M and Mayet S. Psychosocial combined with agonist maintenance treatments versus agonist maintenance treatments alone for treatment of opioid dependence. Cochrane Database of Systematic Reviews 2008, (4 Art. No.: CD004147).

Anderson EN. A healing place: ethnographic notes on a treatment center. Alcoholism Treatment Quarterly, 1992. 9(3–4): 1–21.

Andrews CM, Cao D, Marsh JC and Shin HC. The impact of comprehensive services in substance abuse treatment for women with a history of intimate partner violence. Violence Against Women 2011.17(5):550-67.

Armenakis AA and Bedeian AG. Organizational change: A review of theory and research in the 1990s. Journal of Management, 25, 293-315. 1999.

Ashley OS, Marsden ME and Brady TM. Effectiveness of substance abuse treatment programming for women: a review. American Journal Drug Alcohol Abuse 2003.29(1):19-53.

Australian Bureau of Statistics (ABS). National Survey of Mental Health and Wellbeing: summary of results, Australia, 2012. Canberra: ABS; 2008.

Australian Institute of Health and Welfare. National Drug Strategy Household Survey report Canberra: Australian Institute of Health and Welfare; 2011.

Australian Institute of Health and Welfare. Alcohol and other drug treatment services in Australia 2011-12. Canberra: Australian Institute of Health and Welfare; 2013 and 2014.

Australian Institute of Health and Welfare. National opioid pharmacotherapy statistics annual data collection 2012. Canberra: Australian Institute of Health and Welfare; 2013.

Australian Psychological Society. Evidence-based Psychological Interventions: A Literature Review (Third Edition). Melbourne: The Australian Psychological Society Ltd.; 2010.

Bahr SJ, Masters AL and Taylor BM. What Works in Substance Abuse Treatment Programs for Offenders?; 2012. p. 155-74.

Baker AL, Thornton LK, Hiles S, Hides L and Lubman DI. Psychological interventions for alcohol misuse among people with co-occurring depression or anxiety disorders: A systematic review. Journal of Affective Disorders 2012.139(3):217-29.

Begun AL and Hammond GC. CATCH Court: A Novel Approach to Treatment as Alternative to Incarceration for Women Engaged in Prostitution and Substance Abuse. Journal of Social Work Practice in the Addictions 2012.12(3):328-31.

Bowser BP, Ryan L, Smith CD and Lockett G. Outreach-based drug treatment for sex trading women: the Cal-Pep risk-reduction demonstration project. International Journal of Drug Policy 2008.19(6):492-5.

Brady M. Culture in treatment, culture as treatment: a critical appraisal of developments in addictions programs for Indigenous North Americans and Australians. Social Science and Medicine, 1995 41(11): 1487–1498.

Brady TM and Ashley OS. (Eds.). Women in substance abuse treatment: Results from the Alcohol and Drug Services Study (ADSS) Rockville, MD: Substance Abuse and Mental Health Services Administration, Office of Applied Studies; 2005.

Breen C, Awbery E and Burns L. 2014. Supporting pregnant women who use alcohol or other drugs: a review of the evidence. NDARC, UNSW. Australia.

Breen CL and Burns LA. 2012. Improving services to families affected by FASD: Final report prepared for the Foundation of Alcohol Research and Education, Foundation of Alcohol Research and Education (FARE), Deakin, ACT.

British Columbia Ministry of Health. Service Model and Provincial Standards for Adult Residential Substance Use Services. British Columbia: British Columbia Ministry of Health; 2011.

Briton J. Young people's specialist substance misuse treatment: Exploring the evidence. London: National Treatment Agency for Substance Misuse (NTA); 2009.

Brown CG and Stewart SH. Exploring perceptions of alcohol use as self-medication for depression among women receiving community-based treatment for alcohol problems. Journal of Prevention and Intervention in the Community 2008.35(2):33-47.

Brown JD, Vartivarian S and Alderks CE. Child care in outpatient substance abuse treatment facilities for women: findings from the 2008 National Survey of substance Abuse Treatment Services. Journal of Behavioral Health Services and Research 2011.38(4):478-87.

Brown QL, Cavanaugh CE, Penniman TV and Latimer WW. The impact of homelessness on recent sex trade among pregnant women in drug treatment. Journal of Substance Use 2012.17(3):287-93.

Brown S, Jun MK, Min MO and Tracy EM. Impact of Dual Disorders, Trauma, and Social Support on Quality of Life Among Women in Treatment for Substance Dependence. Journal of Dual Diagnosis 2013.9(1):61-71.

Brown VB and Melchior LA. Women with co-occuring disorders (COD): Treatment settings and service needs. Journal of Psychoactive Drugs 2008.5:365-85.

Browne J and Renzaho, A. Prevention of alcohol and other drug problems in culturally and linguistically diverse communities. Prevention Research Quarterly 2010.13.

Burgess D, Lee R, Tran A and van Ryn, M. 'Effects of Perceived Discrimination on Mental Health and Mental Health Services Utilization Among Gay, Lesbian, Bisexual and Transgender Persons', Journal of LGBT Health Research, 2008 Vol. 3, Iss. 4, pp. 1-14.

Burnette ML, Lucas E, Ilgen M, Frayne SM, Mayo J and Weitlauf JC. Prevalence and health correlates of prostitution among patients entering treatment for substance use disorders. Archives of General Psychiatry 2008.65(3):337-44.

Burnette ML, Schneider R, Ilgen MA and Timko C. Women's past-year prostitution status and receipt of substance abuse treatment services. Psychiatric Services 2008.59(12):1458-61.

Burnette ML, Schneider R, Timko C and Ilgen MA. Impact of substance-use disorder treatment on women involved in prostitution: substance use, mental health, and prostitution one year after treatment. Journal of Studies into Alcohol Drugs 2009.70(1):32-40.

Burns J MC and Thomson N. Summary of Indigenous women's health. Perth, WA: Australian Indigenous HealthInfoNet 2010.

Burns L, Black E, Powers JR, Loxton D, Elliott E, Shakeshaft A, et al. Geographic and maternal characteristics associated with alcohol use in pregnancy. Alcoholism: Clinical and Experimental Research 2011.35(7):1230-7.

Burns L and Breen C. It's time to have the conversation: Understanding the treatment needs of women who are pregnant and alcohol dependent. National Drug and Alcohol Centre University of New South Wales: FARE; 2013.

Burns L, Randall D, Hall WD, Law M, Butler T, et al. Opioid agonist pharmacotherapy in New South Wales from 1985 to 2006: patient characteristics and patterns and predictors of treatment retention. Addiction 2009.104:1363-72.

Burns L, Breen C and Dunlop AJ. 2014. 'Prevention of fetal alcohol spectrum disorders must include maternal treatment.', Medical Journal of Australia, vol. 200, no. 7, pp. 392.

Calabria B, Clifford A, Shakeshaft A, Allan J, Bliss D and Doran C. The acceptability to Aboriginal Australians of a family-based intervention to reduce alcohol-related harms. Drug and Alcohol Review 2013.32(3):328-32.

Campbell CI, Alexander JA and Lemak CH. Organizational determinants of outpatient substance abuse treatment duration in women. Journal of Substance Abuse Treatment 2009.37(1):64-72.

Cavanaugh CE and Latimer WW. Recent sex trade and injection drug use among pregnant opiate and cocaine dependent women in treatment: The significance of psychiatric comorbidity. Addiction Disorders and Their Treatment 2010.9(1):32-40.

Center for Substance Abuse Treatment. A provider's introduction to substance abuse treatment for lesbian, gay, bisexual, and transgender individuals. Rockveill, MD: Substance Abuse and Mental Health Services Administration; 2001.

Center for Substance Abuse Treatment. Addressing the Specific Needs of Women. Rockville, MD: Substance Abuse and Mental Health Services Administration; 2009.

Center for Substance Abuse Treatment. Behavioral Health Services for People Who Are Homeless: A Review of the Literature. Rockville MD: Substance Abuse and Mental Health Services Administration; 2013.

Center for Substance Abuse Treatment. Substance Abuse Among Older Adults. Rockville (MD): Substance Abuse and Mental Health Services Administration; 1998.

Center for Substance Abuse Treatment. Substance Abuse Treatment for Adults in the Criminal Justice System. Rockville MD: Substance Abuse and Mental Health Services Administration; 2005.

Clark TC, Lucassen MF, Bullen P, et al. 2014. The health and well-being of transgender high school students: Results from the New Zealand adolescent health survey (Youth'12). Journal of Adolescent Health 2014. 55: 93-9.

Chen S and Biswas B. Behavioural correlates of dual diagnosis among women in treatment for substance abuse. Mental Health and Substance Use 2012.5(3):217-27.

Cleary M HG, Matheson S and Walter G. Psychosocial treatments for people with co-occurring severe mental illness and substance misuse: systematic review. Journal of Advanced Nursing 2009.65(238-258).

Cleary M HG, Matheson SL, Siegfried N and Walter G. Psychosocial interventions for people with both severe mental illness and substance misuse. Cochrane Database of Systematic Reviews 2008. Issue 1. Art. No.: CD001088.

Colbert AM, Sekula LK, Zoucha R and Cohen SM. Health care needs of women immediately post-incarceration: a mixed methods study. Public Health Nursing 2013.30(5):409-19.

Copeland J. A qualitative study of barriers to formal treatment among women who self-managed change in addictive behaviours. Journal of Substance Abuse Treatment 1997.14(2):183-90.

Copello AG, Velleman RDB and Templeton LJ. Family interventions in the treatment of alcohol and drug problems. Drug Alcohol Rev 2005;24:369 – 385.

Couch M, Pitts M, Mulcare H, Croy S, Mitchell A, Patel S. 2007. Tranznation, A report on the health and well-being of transgender people in Australia and New Zealand, Australian Research Centre in Sex, Health and Society, La Trobe University, Melbourne.

Covington SS, Bloom BE. Gender responsive treatment and services in correctional settings. Women & Therapy 2007.29(3-4):9-33.

Covington SS, Burke C, Keaton S and Norcott C. Evaluation of a trauma-informed and gender-responsive intervention for women in drug treatment. Journal Psychoactive Drugs 2008. Suppl 5:387-98.

Covington SS. Women and Addiction: A Trauma-Informed Approach. Journal of Psychoactive Drugs 2008.S5.

Cusack KJ, Morrissey JP and Ellis AR. Targeting trauma-related interventions and improving outcomes for women with co-occurring disorders. Administrative Policy in Mental Health 2008.35(3):147-58.

Daley KK and Jozica. YSAS Snapshot: Young Women in Youth Alcohol and Other Drug Services. Melbourne, Australia. Youth Support + Advocacy Service; 2013.

Davis KJ and Yonkers KA. Making lemonade out of lemons: a case report and literature review of external pressure as an intervention with pregnant and parenting substance-using women. Journal of Clinical Psychiatry 2012.73(1):51-6.

Davis TA and Ancis J. Look to the relationship: a review of African American women substance users' poor treatment retention and working alliance development. Substance Use & Misuse 2012.47(6):662-72.

Dawe S, Harnett P and Frye S. 2008. Improving outcomes for children living in families with parental substance misuse: What do we know and what should we do? Child Abuse Prevention, 29. Australian Institute of Family Studies.

Dawson A, Jackson D and Cleary M. Mothering on the margins: Homeless women with an SUD and complex mental health co-morbidities. Issues in Mental Health Nursing 2013.34(4):288-93.

Deering KN, Kerr T, Tyndall MW, Montaner JS, Gibson K, Irons L, et al. A peer-led mobile outreach program and increased utilization of detoxification and residential drug treatment among female sex workers who use drugs in a Canadian setting. Drug Alcohol Dependence 2011.113(1):46-54.

Department of Human Services 2002, What's the Difference? Health issues of major concern to gay, lesbian, bisexual, transgender and intersex (GLBTI) Victorians, Department of Human Services, Melbourne.

Dolan K, Salimi S, Nassirimanesh B, Mohsenifar S, Allsop D and Mokri A. Characteristics of Iranian women seeking drug treatment. Journal of Women's Health 2011.20(11):1687-91.

Dolan K, Salimi S, Nassirimanesh B, Mohsenifar S and Mokri A. The establishment of a methadone treatment clinic for women in Tehran, Iran. Journal of Public Health Policy 2011.32(2):219-30.

Drabble L and Eliason MJ. Substance Use Disorders Treatment for Sexual Minority Women. Journal of LGBT Issues in Counseling 2012.6(4):274-92.

Drake RE, O'Neal EL and Wallach MA. A systematic review of psychosocial research on psychosocial interventions for people with co-occurring severe mental and substance use disorders. Journal of Substance Abuse Treatment 2008.34(1):123-38.

Easton H and Matthews R. Evaluation of the Chrysalis Project: Providing accommodation and support for women exiting prostitution. University of Kent; 2012.

Eggleston A, Calhoun PS, Svikis DS, Tuten M, Chisolm MS and Jones HE. Suicidality, aggression, and other treatment considerations among pregnant, substance-dependent women with post traumatic stress disorder. Comprehensive Psychiatry 2009.50(5):415-23.

Eliason MJ, Dibble SL, DeJoseph J and Chinn P. 2009. LGBTQ cultures: What health care professionals need to know about sexual and gender diversity. Philadelphia, PA; 2009.

Ellard J. 2010. Smoking Cessation Interventions: A literature review, AFAO and NAPWA, Sydney Giancas C, Prestage G, Brown G, Triffitt K, Ellard J and Down I. 2013, Experiences of HIV: The seroconversion study annual report 2013, Monograph, the Kirby Institute, the University of New South Wales, Sydney.

Engstrom M, El-Bassel N and Gilbert L. Childhood sexual abuse characteristics, intimate partner violence exposure, and psychological distress among women in methadone treatment. Journal of Substance Abuse Treatment 2012.43(3):366-76.

Engstrom M, Shibusawa T, El-Bassel N and Gilbert L. Age and HIV sexual risk among women in methadone treatment. AIDS Behav 2011.15(1):103-13.

Epperson MW, Khan MR, Miller DP, Perron BE, El-Bassel N and Gilbert L. Assessing criminal justice involvement as an indicator of human immunodeficiency virus risk among women in methadone treatment. Journal of Substance Abuse Treatment 2010.38(4):375-83.

European Monitoring Centre for Drugs and Drug Addiction. Treatment and Care for Older Drug Users. Luxembourg: Publications Office of the European Union; 2010.

Euser SM, Souverein D, Rama Narayana Gowda P, Shekhar Gowda C, Grootendorst D, Ramaiah R, et al. Pragati: an empowerment programme for female sex workers in Bangalore, India. Global Health Action 2012.5:1-11.

Exploring Gender (March 2015) gender centre accessed 12 May 2016 http://www.gendercentre.org.au/resources/fact-sheets/exploring-gender.htm.

Finfgeld-Connett D and Johnson ED. Substance abuse treatment for women who are under correctional supervision in the community: a systematic review of qualitative findings. Issues Mental Health Nursing 2011.32(10):640-8.

Finfgeld-Connett D and Johnson ED. Therapeutic substance abuse treatment for incarcerated women. Clinical Nursing Research 2011.20(4):462-81.

Flaherty I, Donato-Hunt C, Arcuri A and Howard J. Finding the right help: Pathways for culturally diverse clients with cannabis use and mental health issues. Sydney: DAMEC & NCPIC; 2010.

Fowler DN and Faulkner M. Interventions targeting substance abuse among women survivors of intimate partner abuse: a meta-analysis. Journal of Substance Abuse Treatment 2011.41(4):386-98.

Fraser JG, Harris-Britt A, Thakkallapalli EL, Beth K-C and Martin S. Emotional availability and psychosocial correlates among mothers in substance-abuse treatment and their young infants. Infant Mental Health Journal 2010.31(1):1-15.

Gelb K and Rutman D. 2011. Substance Using Women with FASD and FASD Prevention: A Literature Review on Promising Approaches in Substance Use Treatment and Care for Women with FASD. Victoria, BC: University of Victoria.

Graff FS, Morgan TJ, Epstein EE, McCrady BS, Cook SM, Jensen NK, et al. Engagement and retention in outpatient alcoholism treatment for women. American Journal of Addiction 2009.18(4):277-88.

Gray D, Wilson M, Allsop S, Saggers S, Wilkes E and Ober C. Barriers and enablers to the provision of alcohol treatment among Aboriginal Australians: a thematic review of five research projects. Drug and Alcohol Review 2014.

Gray D and Wilkes E. Reducing alcohol and other drug related harm. Canberra: Closing the Gap Clearinghouse; 2010.

Green, C. Gender and use of substance abuse treatment services. Alcohol Research and Health. 2006; 29(1): 55-62.

Green KE and Feinstein BA. Substance use in lesbian, gay, and bisexual populations: an update on empirical research and implications for treatment. Psychology of Addictive Behaviors 2012.26(2):265-78.

Greenfield SF, Brooks AJ, Gordon SM, Green CA, Kropp F, McHugh RK, et al. Substance abuse treatment entry, retention, and outcome in women: a review of the literature. Drug and Alcohol Dependence 2007.86(1):1-21.

Greenfield SF, Pettinati HM, O'Malley S, Randall PK and Randall CL. Gender differences in alcohol treatment: an analysis of outcome from the COMBINE study. Alcoholism: Clinical and Experimental Research 2010.34(10):1803-12.

Greenfield SF and Pirard S. Gender-specific treatment for women with substance use disorders. In: Women and addiction: A comprehensive handbook. New York, NY: Guilford Press; US; 2009. p. 289-306.

Greenfield SF, Rosa C, Putnins SI, Green CA, Brooks AJ, Calsyn DA, et al. Gender research in the National Institute on Drug Abuse National Treatment Clinical Trials Network: a summary of findings. American Journal of Drug and Alcohol Abuse. 2011.37(5):301-12.

Grella CE KM, Warda US, Niv N and Moore AA. . Gender and comorbidity among individuals with opioid use disorders in the NESARC study Addictive Behaviours 2009.346(6-7):498-504.

Grella CE and Greenwell L. Treatment needs and completion of community-based aftercare among substance-abusing women offenders. Women's Health Issues 2007.17(4):244-55.

Grella CE, Needell B, Shi Y and Hser Y-I. Do drug treatment services predict reunification outcomes of mothers and their children in child welfare? Journal of substance abuse treatment 2009.36(3):278-93.

Grella CE and Rodriguez L. Motivation for treatment among women offenders in prison-based treatment and longitudinal outcomes among those who participate in community aftercare.

Journal of Psychoactive Drugs 2011.Suppl 7:58-67.

Grella CE, Scott CK, Foss MA and Dennis ML. Gender similarities and differences in the treatment, relapse, and recovery cycle. Evaluation and Review 2008.32(1):113-37.

Grella CE. From generic to gender-responsive treatment: changes in social policies, treatment services, and outcomes of women in substance abuse treatment. Journal of Psychoactive Drugs 2008. Suppl 5:327-43.

Grella CE. Treatment seeking and utilization among women with substance use disorders. In: Brady KT, Back SE, Greenfield SF, editors. Women & Addiction: A comprehensive handbook. New York, NY: Guilford Press; US; 2009. p. 307-22.

Gruenert S and Tsantefski M. Responding to the needs of children and parents in families experiencing alcohol and other drug problems In: Research Prevention Quarterly, Drug Info, 17 2012.

Guggisberg M. Meeting the needs of women from culturally and linguistically diverse communities. DrugInfo: Drug prevention in culturally and linguistically diverse communities. 2010.

Guarino K, Soares P, Konnath K, Clervil R and Bassuk E. (2009). Trauma-Informed Organizational Toolkit. Rockville, MD: Center for Mental Health Services, Substance Abuse and Mental Health Services Administration, and the Daniels Fund, the National Child Traumatic Stress Network, and the W.K. Kellogg Foundation.

Hall MT, Golder S, Conley CL and Sawning S. Designing programming and interventions for women in the criminal justice system. American Journal of Criminal Justice 2013.38(1):27-50.

Harpaz-Rotem I, Rosenheck RA and Desai R. Residential treatment for homeless female veterans with psychiatric and substance use disorders: effect on 1-year clinical outcomes. Journal of Rehabilitation Research & Development 2011.48(8):891-9.

Harrop EN and Marlatt G. The comorbidity of substance use disorders and eating disorders in women: Prevalence, etiology, and treatment. Addictive Behaviors 2010.35(5):392-8.

Heffner JL BT, Blom, TJ and Anthenelli RM. Gender differences in trauma history and symptoms as predictors of relapse to alcohol and drug use. American Journal of Addiction 2011.20(4):307-11.

Hien DA, Campbell AN, Ruglass LM, Hu MC and Killeen T. The role of alcohol misuse in PTSD outcomes for women in community treatment: a secondary analysis of NIDA's Women and Trauma Study. Drug and Alcohol Dependence 2010.111(1-2):114-9.

Hien DA, Jiang H, Campbell AN, Hu M-C, Miele GM and Cohen LR, et al. Do treatment improvements in PTSD severity affect substance use outcomes? A secondary analysis from a randomized clinical trial in NIDA's Clinical Trials Network. The American Journal of Psychiatry 2010.167(1):95-101.

Hien DA, Morgan-Lopez AA, Campbell ANC, Saavedra LM, Wu E, Cohen L, et al. Attendance and substance use outcomes for the Seeking Safety program: Sometimes less is more. Journal of Consulting and Clinical Psychology 2012.80(1):29-42.

Hien DA, Wells EA, Jiang H, Suarez-Morales L, Campbell AN, Cohen LR, et al. Multisite randomized trial of behavioral interventions for women with co-occurring PTSD and substance use disorders. Journal of Consulting and Clinical Psychology 2009.77(4):607-19.

Hillier L, De Visser R, Kavanagh AM and McNair RP. 2003, 'The Association between Licit and Illicit Drug Use and Sexuality in Young Australian Women', The Medical Journal of Australia, Vol. 179, Iss. 6, pp. 326-327

Hopper E, Bassuk E and Olivet J. Shelter from the Storm: Trauma-Informed Care in Homelessness Services Settings. The Open Health Services and Policy Journal 2010.3:80-100.

Hser YI HD, Teruya C and Anglin MD. Gender differences in treatment outcomes over a three-year period: A path model analysis. Journal of Drug Issues. 2004.34(2):419-39.

Hser YI, Evans E, Huang D and Messina N. Long-term outcomes among drug-dependent mothers treated in women-only versus mixed-gender programs. Journal of Substance Abuse Treatment 2011.41(2):115-23.

Hughes T, McCabe SE, Wilsnack SC, West BT and Boyd CJ. Victimization and substance use disorders in a national sample of heterosexual and sexual minority women and men. Addiction 2010.105(12):2130-40.

Hughes T, Szalacha LA and McNair R. 2010. 'Substance Abuse and Mental Health Disparities: Comparisons across sexual identity groups in a national sample of young Australian women', Social Science and Medicine, Vol. 71, Iss. 4, pp. 824-831.

Hull P, Mao L, Kao S, Edwards B, Prestage G, Zablotska I, de Wit J and Holt M. 2013. Gay Community Periodic Survey Sydney 2013, National Centre in HIV Social Research, the University of New South Wales, Sydney.

Hunt GE, Siegfried N, Morley K, Sitharthan T and Cleary M. Psychosocial interventions for people with both severe mental illness and substance misuse. Cochrane Database Systematic Reviews 2013.10:CD001088.

Hunter DJ and Perkins N. Partnership Working in Public Health. Policy Press. Bristol 2014.

Hutchinson D, Moore EA, Breen C, Burns L and Mattick RP. 2013. 'Alcohol use in pregnancy: prevalence and predictors in the Longitudinal Study of Australian Children.', Drug and Alcohol Review, vol. 32, no. 5, pp. 475 – 482.

Hwang SW, Tolomiczenko G, Kouyoumdjian F and Garner R. Interventions to Improve the Health of the Homeless: A Systematic Review. American Journal of Preventive Medicine 2005.29(4):311.e1-.e9.

Hyde Z, Doherty M, Tilley PJM, McCaul KA, Rooney R, Jancey J. 2014. The First Australian National Trans Mental Health Study: Summary of Results. School of Public Health, Curtin University, Perth, Australia.

Jackson A and Shannon L. Examining barriers to and motivations for substance abuse treatment among pregnant women: does urban-rural residence matter? Women Health 2012.52(6):570-86.

Johnson G, Parkinson S, Tseng, Y and Kuehnle D. Long-term homelessness: Understanding the challenge - 12 months outcomes from the Journey to Social Inclusion pilot program. St Kilda: Sacred Heart Mission; 2011.

Johnson SD, Cottler LB, O'Leary CC and Abdallah AB. The association of trauma and PTSD with the substance use profiles of alcohol- and cocaine-dependent out-of-treatment women. The American Journal on Addictions 2010.19(6):490-5.

Jones HE, Finnegan LP and Kaltenbach K. Methadone and Buprenorphine for the Management of Opioid Dependence in Pregnancy. Drugs & Alcohol Today 2012.72(6):747-57.

Jones HE, O'Grady KE and Tuten M. Reinforcement-based treatment improves the maternal treatment and neonatal outcomes of pregnant patients enrolled in comprehensive care treatment. American Journal of Addiction 2011.20(3):196-204.

Kay A, Taylor TE, Barthwell AG, Wichelecki J and Leopold V. Substance use and women's health. Journal of Addictive Diseases 2010.29(2):139-63.

Keegan J, Parva M, Finnegan M, Gerson A and Belden M. Addiction in pregnancy. Journal of Addictive Diseases 2010.29(2):175-91.

Kelly SM, Schwartz RP, O'Grady KE, Mitchell SG, Reisinger HS, Peterson JA, et al. Gender Differences Among In- and Out-of-Treatment Opioid-Addicted Individuals. American Journal of Drug Alcohol Abuse 2009.35(1):38-42.

Kennare R, Heard A and Chan A. Substance use during pregnancy: risk factors and obstetric and perinatal outcomes in South Australia. Australian and New Zealand Journal of Obstetrics and Gynaecology 2005.45:220-5.

Klein AA and Slaymaker VJ. 12-Step Involvement and Treatment Outcomes Among Young Women With Substance Use Disorders. Alcoholism Treatment Quarterly 2011.29(3):204-18.

Koehler JA, Humphreys DK, Akoensi TD, Sánchez de Ribera O and Lösel F. A systematic review and metaanalysis on the effects of European drug treatment programmes on reoffending. Psychology, Crime & Law 2013.20(6):584-602.

Kratzmann M ME, Ware J, Banach L, Ward J, Ryan J, Sutton L, Griffiths P and Saunders M. Injecting drug use and associated harms among Aboriginal Australians. Canberra: Australian National Council on Drugs; 2011.

Lea T, Reynolds R and de Wit J. Alcohol and other drug use, club drug dependence and treatment seeking among lesbian, gay and bisexual young people in Sydney. Drug Alcohol Rev 2013.32(3):303-11.

Lee KK, Harrison K, Mills K and Conigrave KM. Needs of Aboriginal Australian women with comorbid mental and alcohol and other drug use disorders. Drug and Alcohol Review 2014.

Lee SK, Sulaiman-Hill CM and Thompson SC. Alcohol experiences and concerns of newly arrived migrant women. 2014 Sage Open 4(2).

Lee KS, Dawson A and Conigrave KM. The role of an Aboriginal women's group in meeting the high needs of clients attending outpatient alcohol and other drug treatment. Drug and Alcohol Review 2013.

Lester BM, Twomey JE. Treatment of substance abuse during pregnancy. Womens Health 2008.4:67-77.

Li Q, Li X and Stanton B. Alcohol use among female sex workers and male clients: An integrative review of global literature. Alcohol and Alcoholism 2010.45(2):188-99.

Linehan MM, Dimeff LA, Reynolds SK, Comtois KA, Welch SS, Heagerty P, et al. Dialectical behavior therapy versus comprehensive validation therapy plus 12-step for the treatment of opioid dependent women meeting criteria for borderline personality disorder. Drug and Alcohol Dependence 2002.67(1):13-26.

Linehan MM, Schmidt H, 3rd, Dimeff LA, Craft JC, Kanter J and Comtois KA. Dialectical behavior therapy for patients with borderline personality disorder and drug-dependence. American Journal of Addiction 1999.8(4):279-92.

Lipsky S, Krupski A, Roy-Byrne P, Huber A, Lucenko BA and Mancuso D. Impact of sexual orientation and co-occurring disorders on chemical dependency treatment outcomes. Journal of Studies on Alcohol and Drugs. 2012.73(3):401-12.

Loxley W, Adams K, Women, drug use and crime: findings from the Drug Use Monitoring in Australia program. Research and public policy Series 99. Canberra: Australian Institute of Criminology, 2009.

Lui ST M and Smith EJ. Psychosocial interventions for women enrolled in alcohol treatment during pregnancy. Cochrane Database of Systematic Reviews 2008. (3 Art. No.: CD006753).

Lundgren LM, Schilling RF, Fitzgerald T, Davis K and Amodeo M. Parental status of women injection drug users and entry to methadone maintenance. Substance Use & Misuse 2003.38(8):1109-31.

Magee C and Huriaux E. Ladies' night: Evaluating a drop-in programme for homeless and marginally housed women in San Francisco's mission district. International Journal of Drug Policy 2008.19(2):113-21.

Maher L and Hudson SL. Women in the Drug Economy: A Metasynthesis of the Qualitative Literature. Journal of Drug Issues 2007 37:805-826.

Massey SH, Lieberman DZ, Reiss D, Leve LD, Shaw DS and Neiderhiser JM. Association of clinical characteristics and cessation of tobacco, alcohol, and illicit drug use during pregnancy. American Journal on Addictions 2011.20(2):143-50.

Massey SH, Neiderhiser JM, Shaw DS, Leve LD, Ganiban JM and Reiss D. Maternal self concept as a provider and cessation of substance use during pregnancy. Addictive Behaviors 2012.37(8):956-61.

Matheson FI, Doherty S and Grant BA. Community-based aftercare and return to custody in a national sample of substance-abusing women offenders. American Journal of Public Health 2011.101(6):1126-32.

McCabe JE and Arndt S. Demographic and substance abuse trends among pregnant and non-pregnant women: eleven years of treatment admission data. Maternal and Child Health Journal 2012.16(8):1696-702.

McCabe SE, West BT, Hughes TL and Boyd CJ. Sexual orientation and substance abuse treatment utilization in the United States: results from a national survey. Journal of Substance Abuse Treatment 2013.44(1):4-12.

McCormick RM. Aboriginal traditions in the treatment of substance abuse. Canadian Journal of Counselling, 2000. 34(1): 25–32.

McCrady BS, Epstein EE, Cook S, Jensen N and Hildebrandt T. A randomized trial of individual and couple behavioral alcohol treatment for women. Journal of Consulting Clinical Psychology 2009.77(2):243-56.

McDonald, D. Case management in the substance abuse field: insights from the research literature. A report prepared for the Drug & Alcohol Policy Unit ACT Health. 2005.

Messina N, Burdon W and Prendergast M. Prison-based treatment for drug-dependent women offenders: treatment versus no treatment. Journal of Psychoactive Drugs 2006. Suppl 3:333-43.

Messina N, Calhoun S and Warda U. Gender-responsive drug court treatment: A randomized controlled trial. Criminal Justice and Behavior. 2012.39(12):1539-58.

Meyers RJ, Roozen HG and Smith JE. The Community Reinforcement Approach: An Update of the Evidence. Alcohol Research & Health 2011.33(4):380-8.

Milligan K, Niccols A, Sword W, Thabane L, Henderson J, Smith A, et al. Maternal substance use and integrated treatment programs for women with substance abuse issues and their children: a meta-analysis. Substance Abuse Treatment, Prevention and Policy 2010.5:21.

Milligan K, Niccols A, Sword W, Thabane L, Henderson J and Smith A. Birth outcomes for infants born to women participating in integrated substance abuse treatment programs: A meta-analytic review. Addiction Research & Theory 2011.19(6):542-55.

Milligan K, Niccols A, Sword W, Thabane L, Henderson J and Smith A. Length of stay and treatment completion for mothers with substance abuse issues in integrated treatment programmes. Drugs: Education, Prevention, and Policy 2011.18(3):219-27.

Mills KD, M, Proudfoot H, Sannibale C, Teeson M, Mattick R and Burns, L. Guidelines on the management of co-occurring alcohol and other drug and mental health conditions in alcohol and other drug treatment settings. Sydney: National Drug and Alcohol Research Centre, University of NSW; 2009.

Ministerial Council on Drug Strategy. National drug strategy 2010-2015: a framework for action on alcohol, tobacco and other drugs. Canberra: Commonwealth of Australia; 2011.

Minozzi S, Amato L, Vecchi S and Davoli M. Maintenance agonist treatments for opiate dependent pregnant women. Cochrane Database Systems Review 2008. (2):CD006318.

Mitchell MM, Severtson SG and Latimer WW. Pregnancy and race/ethnicity as predictors of motivation for drug treatment. American Journal of Drug and Alcohol Abuse 2008.34(4):397-404.

Mooney Somers J, Deacon RM, Comfort J, Richters J and Parkhill N. 2013, Women in Contact with the Sydney Gay and Lesbian Community: Report of the Sydney Women and Sexual Health (SWASH) Survey 2006, 2008, 2010 and 2012, ACON and Centre for Values, Ethics and the Law in Medicine (VELiM), University of Sydney, Sydney.

Morrissey JP, Jackson EW, Ellis AR, Amaro H, Brown VB and Najavits LM. Twelve-month outcomes of trauma-informed interventions for women with co-occurring disorders. Psychiatric Services 2005.56(10):1213-22.

Myers B, Louw J and Pasche S. Gender differences in barriers to alcohol and other drug treatment in Cape Town, South Africa. African Journal of Psychiatry (Johannesburg) 2011.14(2):146-53.

Network of Alcohol and other Drugs Agencies (NADA) Tools for Change: A new way of working with families and carers, NADA 2009

Network of Alcohol and other Drugs Agencies (NADA) Complex Needs Capable: A Practice Resource for Drug and Alcohol Services NADA 2013

Network of Alcohol and other Drugs Agencies (NADA) Working with Diversity in Alcohol and other Drug Settings NADA 2014

Najavits LM. Psychotherapies for trauma and substance abuse in women: review and policy implications. Trauma Violence Abuse 2009.10(3):290-8.

Nagel T, Robinson G, Condon J and Trauer T. Approach to treatment of mental illness and substance dependence in remote Indigenous communities: results of a mixed methods study. Australian Journal of Rural Health, 2009. 17(4): 174–182.

National Institute on Drug Abuse. Principles of Drug Addiction Treatment: A Research Based Guide (Second Edition). Services USDoHaH. National Institutes of Health; 2009.

Niccols A MK, Sword W, Thabane L, Henderson J and Smith A. Integrated programs for mothers with substance abuse issues: A systematic review of studies reporting on parenting outcomes. Harm Reduction Journal 2012.9(14).

Niccols A, Dell CA and Clarke S. Treatment Issues for Aboriginal Mothers with Substance Use Problems and Their Children. International Journal of Mental Health and Addiction 2010.8(2):320-35.

Niccols A, Milligan K, Smith A, Sword W, Thabane L and Henderson J. Integrated programs for mothers with substance abuse issues and their children: a systematic review of studies reporting on child outcomes. Child Abuse and Neglect 2012.36(4):308-22.

NSW Community Services (2010). Working with parental substance misuse. Research to Practice Notes. February 2010.

NSW Department of Health. National Clinical Guidelines for the Management of Drug Use during Pregnancy, Birth and the Early Development Years of the Newborn. Sydney: NSW Department of Health; 2006.

NSW Family and Community Services (2014). Shining a light on good practice in NSW in 2014: stories from child protection and out-of-home care.

Nuttbrock LA, Rosenblum A, Magura S, Villano C and Wallace J. Linking female sex workers with substance abuse treatment. Journal of Substance Abuse Treatment 2004.27(3):233-9.

Ondersma SJ, Winhusen T and Lewis DF. External pressure, motivation, and treatment outcome among pregnant substance-using women. Drug and Alcohol Dependence. 2010.107(2-3):149-53.

Osterling KL and Austin MJ. Substance abuse interventions for parents involved in the child welfare system: evidence and implications. Journal of Evidence Based Social Work 2008.5(1-2):157-89.

Pajulo M, Pyykkonen N, Kalland M, Sinkkonen J, Helenius H, Punamaki RL, et al. Substance-Abusing Mothers in Residential Treatment with Their Babies: Importance of Pre- and Postnatal Maternal Reflective Functioning. Infant Mental Health J 2012.33(1):70-81.

Pajulo M, Pyykkonen N, Kalland M, Sinkkonen J, Helenius H and Punamaki RL. Substance abusing mothers in residential treatment with their babies: postnatal psychiatric symptomatology and its association with mother-child relationship and later need for child protection actions. Nordic Journal of Psychiatry 2011.65(1):65-73.

Panchanadeswaran S and Jayasundara D. Experiences of Drug Use and Parenting among Women in Substance Abuse Treatment: An Exploratory Study. Journal of Human Behavior in the Social Environment 2012.22(8):971-87.

Pinkham S and Malinowska-Sempruch K. Women, Harm Reduction, and HIV. New York: International Harm Reduction Development Program of the Open Society Institute. 2007.

Prendergast ML, Messina NP, Hall EA and Warda US. The relative effectiveness of women-only and mixed-gender treatment for substance-abusing women. Journal of Substance Abuse Treatment 2011.40(4):336-48.

Rasmussen C, Kully-Martens K, Denys K, Badry D, Henneveld D, Wyper K, et al. The effectiveness of a community-based intervention program for women at-risk for giving birth to a child with Fetal Alcohol Spectrum Disorder (FASD). Community Mental Health Journal 2012.48(1):12-21.

Ranzijn R, McConnochie K and Nolan W. Psychology and Indigenous Australian, Foundations of Culture Competence, Palgrave Macmillan: South Yarra, Victoria. 2010.

Resko SM and Mendoza NS. Early Attrition From Treatment Among Women With Co-occurring Substance Use Disorders and PTSD. Journal of Social Work Practice in the Addictions 2012.12(4):348-69.

Ritter, Matthew-Simons and Carragher 2012:67 Prevalence of and interventions for mental health and Alcohol and Other Drug Problems amongst the Gay, Lesbian, Bisexual and Transgender Community: A review of the literature, Drug Policy Modelling Program, Monograph Series no. 23 NDARC.

Roarty L and Saggers S. Evaluation of services to culturally and linguistically diverse (CALD) women with comorbid mental health and drug and alcohol issues. Perth: NDRI; 2009.

Rosen D, Hunsaker A, Albert SM, Cornelius JR and Reynolds CF. Characteristics and consequences of heroin use among older adults in the United States: A review of the literature, treatment implications, and recommendations for further research. Addictive Behaviors 2011.36(4):279-85.

Rösner S H-HA, Leucht S, Lehert P, Vecchi S and Soyka M. Acamprosate for alcohol dependence. Cochrane Database of Systematic Reviews 2010. (Issue 9. Art. No.: CD004332.).

Rösner S H-HA, Leucht S, Vecchi S, Srisurapanont M and Soyka M. Opioid antagonists for alcohol dependence. Cochrane Database of Systematic Reviews 2010. (Issue 12. Art. No.: CD001867).

Rowan-Szal GA, Joe GW, Simpson DD, Greener JM and Vance J. During-treatment Outcomes among Female Methamphetamine-Using Offenders in Prison-based Treatments. Journal of Offender Rehabilitation 2009.48(5):388-401.

Rowe R. Choosing the right mix: lessons on culturally relevant treatment from the evaluation of the Drug and Alcohol Multicultural Education Centre's Counselling Service. Mental Health and Substance Use 2013.7(2):92-101.

Roxburgh A, Degenhardt L, Copeland J and Larance B. Drug dependence and associated risks among female street-based sex workers in the greater Sydney area, Australia. Substance Use and Misuse 2008.43(8-9):1202-17.

Roxburgh A, Degenhardt L and Copeland J. Posttraumatic stress disorder among female street-based sex workers in the greater Sydney area, Australia. British Medical College of Psychiatry 2006.6(1):24.

Royal College of Psychiatrists. Our Invisible Addicts: First Report of the Older Person's Substance Misuse Working Group of the Royal College of Psychiatrists. London: Royal College of Psychiatrists; 2011.

Ruglass LM, Miele GM, Hien DA, Campbell AN, Hu MC, Caldeira N, et al. Helping alliance, retention, and treatment outcomes: a secondary analysis from the NIDA Clinical Trials Network Women and Trauma Study. Substance Use Misuse 2012.47(6):695-707.

Sacks JY, McKendrick K and Banks S. The impact of early trauma and abuse on residential substance abuse treatment outcomes for women. Journal of Substance Abuse Treatment 2008.34(1):90-100.

Sacks JY, McKendrick K and Hamilton Z. A randomized clinical trial of a therapeutic community treatment for female inmates: outcomes at 6 and 12 months after prison release. Journal of Addictive Diseases 2012.31(3):258-69.

Sallmann J. "Going Hand-in-Hand": Connections Between Women's Prostitution and Substance Use. Journal of Social Work Practice in the Addictions 2010.10(2):115-38.

Sallmann J. Living With Stigma: Women's Experiences of Prostitution and Substance Use. Affilia 2010.25(2):146-59.

SAMHSA 2006 (Substance Abuse and Mental Health Services Administration). Developing a Stigma Reduction Initiative. SAMHSA Pub No. SMA-4176. Rockville MD. Center for Mental Health Services, Substance Abuse and Mental Health Services Administration.

Scott CK, Dennis ML. The first 90 days following release from jail: findings from the Recovery Management Checkups for Women Offenders (RMCWO) experiment. Drug and Alcohol Dependence 2012.125(1-2):110-8.

Seib C, Fischer J and Najman J. The health of female sex workers from three industry sectors in Queensland. Social Science and Medicine. 68(3) 2008: 473-478.

Senreich E. A Comparison of Perceptions, Reported Abstinence, and Completion Rates of Gay, Lesbian, Bisexual, and Heterosexual Clients in Substance Abuse Treatment. Journal of Gay & Lesbian Mental Health 2009.13(3):145-69.

Sevelius JM, Reznick OG, Hart S and Schwarcz S. Informing interventions: The Importance of contextual factors in the prediction of sexual risk behaviors among transgender women. AIDS Education and Prevention 2009.21(2):113-27.

Shannon K, Kerr T, Allinott S, Chettiar K, Shoveller and Tyndall MW. Social and structural violence and power relations in mitigating HIV risk of drug-using women in survival sex work Social Science and Medicine Volume 66 (4) 2008:911–921.

Shimmin C. 2009 Understanding stigma through a gender lens. Canadian Women's Health Network, 11:2.

Skinner N. Organisational Change. In Skinner N, Roche AM, O'Connor J, Pollard Y and Todd C.(Eds.), Workforce Development TIPS (Theory Into Practice Strategies): A Resource Kit for the Alcohol and Other Drugs Field. National Centre for Education and Training on Addiction (NCETA), Flinders University, Adelaide, Australia 2005.

Simpson M and McNulty J. Different needs: women's drug use and treatment in the UK. International Journal of Drug Policy 2008.19(2):169-75.

Slaymaker VJ and Owen P. Alcohol and other drug dependence severity among older adults in treatment: Measuring characteristics and outcomes. Alcoholism Treatment Quarterly 2008.26(3):259-73.

Slesnick N and Erdem G. Intervention for homeless, substance abusing mothers: findings from a non-randomized pilot. Behavioral Medicine 2012.38(2):36-48.

Smith EJ, Lui S and Terplan M. Pharmacologic interventions for pregnant women enrolled in alcohol treatment. Cochrane Database Systems Review 2009.3(3):CD007361.

Smith IE. Prevention and Intervention Programs for Pregnant Women Who Abuse Substances. In: Straussner SLA, Fewell CH, editors. Children of Substance-Abusing Parents: Dynamics and Treatment. New York: Springer Publishing Company; 2011. p. 171-92.

Speirs V, Johnson M and Jirojwong S. A systematic review of interventions for homeless women. Journal of Clinical Nursing 2013.22(7-8):1080-93.

Spooner C and Hetherington K. Social Determinants of Drug Use. Sydney: National Drug and Alcohol Research Centre; 2004.

Storbjörk J. Gender differences in substance use, problems, social situation and treatment experiences among clients entering addiction treatment in Stockholm. Nordic Studies on Alcohol and Drugs 2011.28(3):185-209.

Suchman NE, DeCoste C, Castiglioni N, McMahon TJ, Rounsaville B and Mayes L. The Mothers and Toddlers Program, an attachment-based parenting intervention for substance using women: post-treatment results from a randomized clinical pilot. Attachment and Human Development 2010.12(5):483-504.

Suchman NE, Decoste C, McMahon TJ, Rounsaville B, Mayes L. The Mothers and Toddlers program, an attachment-based parenting intervention for substance-using women: Results at 6-week follow-up in a randomized clinical pilot. Infant Mental Health Journal 2011.32(4):427-49.

Swift W, Copeland J, Hall W. Characteristics of women with alcohol and other drug problems: findings of an Australian national survey. Addiction 1996.91(8):1141-50.

Tait RJ, French DJ, Burns R, Anstey KJ. Alcohol use and depression from middle age to the oldest old: gender is more important than age. International Psychogeriatrics 2012.24(8):1275-83.

Taplin S, Mattick RP. Child Protection and Mothers in Substance Abuse Treatment Sydney, NSW: National Drug and Alcohol Research Centre; 2011.

Taylor K, Thompson S, Davis R. Delivering culturally appropriate residential rehabilitation for urban Indigenous Australians: a review of the challenges and opportunities. Australian and New Zealand Journal of Public Health 2010.34 Suppl 1:S36-40.

Taylor L, Hutchinson D, Rapee R, Burns L, Stephens C and Haber PS. Clinical features and correlates of outcomes for high-risk, marginalized mothers and newborn infants engaged with a specialist perinatal and family drug health service. Obstetrics and Gynaecology International 2012.2012:867265.

Taylor O. Barriers to Treatment for Women with Substance Use Disorders. Journal of Human Behavior in the Social Environment 2010.20(3):393-409.

Teasdale K, Conigrave K, Kiel K, Freeburn B, Long G and Becker K. Improving services for prevention and treatment of substance misuse for Aboriginal communities in a Sydney Area Health Service Drug and Alcohol Review 2008.27(152-159).

Terplan M, Garrett J, Hartmann K. Gestational age at enrolment and continued substance use among pregnant women in drug treatment. Journal of Addictive Disease 2009.28(2):103-12.

Terplan M and Lui S. Psychosocial interventions for pregnant women in outpatient illicit drug treatment programs compared to other interventions. John Wiley and Sons, Ltd; 2007.

Terplan M, McNamara EJ and Chisolm MS. Pregnant and non-pregnant women with substance use disorders: the gap between treatment need and receipt. .Journal of Addictive Disease 2012.31(4):342-9.

Terrell, M.D. Ethnocultural factors and substance abuse: toward culturally sensitive treatment models. Psychology of Addictive Behaviors, 1993 7(3): 162–167.

The Jean Tweed Centre. Trauma Matters: Guidelines for Trauma-Informed Practices in Women's Substance Use Services. The Jean Tweed Centre, Ontario. 2013.

Tiet QQ MB. Treatment for patients with dual diagnosis: a review. Alcoholism: Clinical and Experimental Research in Nursing & Health 2007.31(4):513-36.

Trifonoff A, Duraisingam V, Roche AM and Pidd K. 2010. Taking First Steps: What family sensitive practice means for alcohol and other drug workers, survey report. NCETA SA.

Tripodi SJ, Bledsoe SE, Kim JS and Bender K. Effects of Correctional-Based Programs for Female Inmates: A Systematic Review. Research on Social Work Practice 2011.21(1):15-31.

Tripodi SJ and Pettus-Davis C. Histories of childhood victimization and subsequent mental health problems, substance use, and sexual victimization for a sample of incarcerated women in the US. International Journal of Law and Psychiatry 2013.36(1):30-40.

Tuchman E. Women and addiction: the importance of gender issues in substance abuse research. Journal of Addictive Diseases 2010.29(2):127-38.

Turnbull C and Osborn DA. Home visits during pregnancy and after birth for women with an alcohol or drug problem. John Wiley & Sons, Ltd; 2012.

Understanding gender (n.d) gender spectrum accessed 12 May 2016, https://www.genderspectrum.org/quick-links/understanding-gender.

Wechsberg WM, Wu LT, Zule WA, Parry CD, Browne FA, Luseno WK, et al. Substance abuse, treatment needs and access among female sex workers and non-sex workers in Pretoria, South Africa. Substance Abuse Treatment, Prevention and Policy2009.4:11.

Whitaker T, Ryan P and Cox G. Stigmatization among drug-using sex workers accessing support services in Dublin. Qualitative Health Research 2011.21(8):1086-100.

Willis K. and Rushforth C. The Female Criminal: An Overview of Women's Drug Use and Offending Behaviour. Trends & Issues in Crime and Criminal Justice 2003:264.

Wilson EC, Garofalo R, Harris RD, Herrick A, Martinez M, Martinez J and Belzer M. Transgender female youth and sex work: HIV risk and a comparison of life factors related to engagement in sex work. AIDS and Behavior. 2009. p. 902-13.

Winhusen T, Kropp F, Babcock D, Hague D, Erickson SJ, Renz C, et al. Motivational enhancement therapy to improve treatment utilization and outcome in pregnant substance users. Journal of Substance Abuse Treatment 2008.35(2):161-73.

Winklbaur B, Kopf N, Ebner N, Jung E, Thau K and Fischer G. Treating pregnant women dependent on opioids is not the same as treating pregnancy and opioid dependence: a knowledge synthesis for better treatment for women and neonates. Addiction 2008.103(9):1429-40.

Witkiewitz K, Greenfield BL and Bowen S. Mindfulness-based relapse prevention with racial and ethnic minority women. Addictive Behaviors 2013.38(12):2821-4.

Wu LT and Blazer DG. Illicit and nonmedical drug use among older adults: A review. Journal of Aging and Health 2011.23(3):481-504.

Yeom HS. Gender differences in treatment outcomes among participants in a mixed-gender substance abuse aftercare program. Social Work in Public Health 2011.26(6):557-76.

Yonkers KA, Forray A, Howell HB, Gotman N, Kershaw T, Rounsaville BJ, et al. Motivational enhancement therapy coupled with cognitive behavioral therapy versus brief advice: a randomized trial for treatment of hazardous substance use in pregnancy and after delivery. General Hospital Psychiatry 2012.34(5):439-49.

Young JL and Martin PR. Treatment of opioid dependence in the setting of pregnancy. Psychiatric Clinics of North America 2012.35(2):441-60.

# NOTES

NOTES



